arXiv:0807.0973v1 [math.DG] 7 Jul 2008

AN END-TO-END CONSTRUCTION FOR SINGLY PERIODIC MINIMAL SURFACES

L. HAUSWIRTH, F. MORABITO, AND M. M. RODRÍGUEZ

Abstract. We show the existence of various families of properly embedded singly periodic minimal surfaces in $\mathbb{R}^3$ with finite arbitrary genus and Scherk type ends in the quotient. The proof of our results is based on the gluing of small perturbations of pieces of already known minimal surfaces.

1. Introduction

Besides the plane and the helicoid, the first singly periodic minimal surface was discovered by Scherk [32] in 1835. This surface, known as *Scherk's second surface*, is a properly embedded minimal surface in $\mathbb{R}^3$ invariant by one translation T we can assume along the x_2-axis, and can be seen as the desingularization of two perpendicular planes P_1 and P_2 containing the x_2-axis. We assume P_1, P_2 are symmetric with respect to the planes $\{x_1 = 0\}$ and $\{x_3 = 0\}$. By changing the angle between P_1, P_2 we obtain a 1-parameter family of properly embedded singly periodic minimal surfaces, we will refer to as *Scherk surfaces*. In the quotient $\mathbb{R}^3/T$ by its shortest translation T, each Scherk surface has genus zero and four ends asymptotic to flat annuli contained in $P_1/T, P_2/T$. Such ends are called Scherk-type ends.

In 1982, C. Costa [2, 3] discovered a genus one minimal surface with three embedded ends: one top catenoidal end, one middle planar end and one bottom catenoidal end. D. Hoffmann and W.H. Meeks [8, 9, 10] proved the global embeddedness for this Costa example, and generalized it for higher genus. For each $k \geqslant 1$, Costa-Hoffman-Meeks surface M_k (we will abbreviate by saying *CHM example*) is a properly embedded minimal surface of genus k and three ends: two catenoidal ones and one middle planar end.

F. Martin and V. Ramos Batista [18] have recently constructed a properly embedded singly periodic minimal example which has genus one and six Scherk-type ends in the quotient $\mathbb{R}^3/T$, called *Scherk-Costa surface*, based on Costa surface (from now on, T will denote a translation in the x_2-direction). Roughly speaking, they remove each end of Costa surface (asymptotic to a catenoid or a plane) and replace it by two Scherk-type ends. In this paper we obtain surfaces in the same spirit as Martin and Ramos Batista's one, but with a completely different method. We construct properly embedded singly periodic minimal surfaces with genus $k \geqslant 1$ and six Scherk-type ends in the quotient $\mathbb{R}^3/T$, by gluing (in an analytic way) a compact piece of M_k to two halves of a Scherk surface at the top and bottom catenoidal ends, and one flat horizontal annulus P/T with a disk removed at the middle planar end.

1991 *Mathematics Subject Classification.* 53A10, 49Q05.

Third author research activity is partially supported by grants from Région Ile-de-France and a MEC/FEDER grant no. MTM2007-61775.

Theorem 1.1. *Let T denote a translation in the x_2-direction. For each $k \geqslant 1$, there exists a 1-parameter family of properly embedded singly periodic minimal surfaces in $\mathbb{R}^3$ invariant by T whose quotient in $\mathbb{R}^3/T$ has genus k and six Scherk-type ends.*

V. Ramos Batista [29] constructed a singly periodic Costa minimal surface with two catenoidal ends and two Scherk-type middle end, which has genus one in the quotient $\mathbb{R}^3/T$. This example is not embedded outside a slab in $\mathbb{R}^3/T$ which contains the topology of the surface. We observe that the surface we obtain by gluing a compact piece of M_1 (Costa surface) at its middle planar end to a flat horizontal annulus with a disk removed has the same properties as Ramos Batista's one.

In 1988, H. Karcher [13, 14] defined a family of properly embedded doubly periodic minimal surfaces, called *toroidal halfplane layers*, which has genus one and four horizontal Scherk-type ends in the quotient. In 1989, W.H. Meeks and H. Rosenberg [21] developed a general theory for doubly periodic minimal surfaces having finite topology in the quotient, and used an approach of minimax type to obtain the existence of a family of properly embedded doubly periodic minimal surfaces, also with genus one and four horizontal Scherk-type ends in the quotient. These Karcher's and Meeks and Rosenberg's surfaces have been generalized in [30], constructing a 3-parameter family $\mathcal{K} = \{M_{\sigma,\alpha,\beta}\}_{\sigma,\alpha,\beta}$ of surfaces, called *KMR examples* (sometimes, they are also referred in the literature as toroidal halfplane layers). Such examples have been classified by J. Pérez, M.M. Rodríguez and M. Traizet [27] as the only properly embedded doubly periodic minimal surfaces with genus one and finitely many parallel (Scherk-type) ends in the quotient. Each $M_{\sigma,\alpha,\beta}$ is invariant by a horizontal translation T (by the period vector at its ends) and a non horizontal one $\widetilde{T}$. We denote by $\widetilde{M}_{\sigma,\alpha,\beta}$ the lifting of $M_{\sigma,\alpha,\beta}$ to $\mathbb{R}^3/T$, which has genus zero, infinitely many horizontal Scherk-type ends, and two limit ends.

In 1992, F.S. Wei [33] added a handle to a KMR example $M_{\sigma,0,0}$ in a periodic way, obtaining a properly embedded doubly periodic minimal surface invariant under reflection in three orthogonal planes, which has genus two and four horizontal Scherk-type ends in the quotient. Some years later, W. Rossman, E.C. Thayer and M. Wolgemuth [31] added a different type of handle to a KMR example $M_{\sigma,0,0}$, also in a periodic way, obtaining a different minimal surfaces with the same properties as Wei's one. They also added two handles to a KMR example, getting doubly periodic examples of genus three in the quotient. L. Mazet and M. Traizet [20] have recently added $N \geqslant 1$ handles to a KMR example $M_{\sigma,0,0}$, obtaining a genus N properly embedded minimal surface in $\mathbb{R}^3/T$ with an infinite number of horizontal Scherk-type ends and two limit ends. The idea of the construction is to connect N periods of the doubly periodic example of Wei with two halves KMR example. However they only control the asymptotic behavior in their construction. They have also constructed a properly embedded minimal surface in $\mathbb{R}^3/T$ with infinite genus, adding handles in a quasi-periodic way to a KMR example.

L. Hauswirth and F. Pacard [6] have constructed higher genus Riemann minimal examples in $\mathbb{R}^3$, by gluing two halves of a Riemann minimal example with the intersection of a conveniently chosen CHM surface with a slab. We follow their ideas to generalize Mazet and Traizet's examples by constructing higher genus KMR examples: we construct properly embedded singly periodic minimal examples whose quotient in $\mathbb{R}^3/T$ has arbitrary finite genus, infinitely many horizontal Scherk-type ends and two limit ends. More precisely, we glue a compact piece of a slightly deformed CHM example M_k with tilted catenoidal ends, to two halves of a KMR example $M_{\sigma,\alpha,0}$ or $M_{\sigma,0,\beta}$ (see Figure 1) and a periodic horizontal flat annuli with a disk removed.



FIGURE 1. A sketch of half a KMR example $M_{\sigma,0,0}$ glued to a compact piece of Costa surface.

Theorem 1.2. *Let T denote a translation in the x_2-direction. For each $k \geqslant 1$, there exist two 1-parameter families $\mathcal{K}_1, \mathcal{K}_2$ of properly embedded singly periodic minimal surfaces in $\mathbb{R}^3$ whose quotient in $\mathbb{R}^3/T$ has genus k, infinitely many horizontal Scherk-type ends and two limit ends. The surfaces in $\mathcal{K}_1$ have a vertical plane of symmetry orthogonal to the x_1-axis, and the surfaces in $\mathcal{K}_2$ have a vertical plane of symmetry orthogonal to the x_2-axis.*

L. Mazet, M. Traizet and M. Rodríguez [19] have recently constructed saddle towers with infinitely many ends: they are non-periodic properly embedded minimal surfaces in $\mathbb{R}^3/T$ with infinitely many ends and one limit end. In the present paper, we construct (non-periodic) properly embedded minimal surfaces in $\mathbb{R}^3/T$ with arbitrary finite genus $k \geqslant 0$, infinitely many ends and one limit end. With this aim, we glue half a Scherk example with half a KMR example, in the case $k = 0$; and, when $k \geqslant 1$, we glue a compact piece of the CHM example M_k to half a Scherk surface (at the top catenoidal end of M_k), a periodic horizontal flat annuli with a disk removed (at the middle planar end) and half a KMR example (at the bottom catenoidal end), see Figure 2.

Theorem 1.3. *Let T denote a translation in the x_2-direction. For each $k \geqslant 0$, there exists a 1-parameter family of properly embedded singly periodic minimal surfaces in $\mathbb{R}^3$ whose quotient in $\mathbb{R}^3/T$ has genus k, infinitely many horizontal Scherk-type ends and one limit end.*

The family of KMR examples is a three parameter family which contains two subfamilies whose surfaces have a vertical plane of symmetry. In the construction of examples satisfying theorems 1.2 and 1.3, we need to have at least one vertical plane of symmetry in order to control the kernel of the Jacobi operator on each glued piece. F. Morabito [24] has recently proved there is a bounded Jacobi field which does not come from isometries of $\mathbb{R}^3$ on the CHM bent surface. For this reason, we are not able to produce a 3-parameter family of KMR examples with higher genus in theorem 1.2.

The paper is organized as follows: In section 2 we briefly describe the CHM examples M_k and obtain, for each genus k, a 1-parameter family of surfaces $M_k(\xi)$ by bending the catenoidal ends of $M_k = M_k(0)$ keeping a vertical plane of symmetry. This is used to prescribe the flux of the deformed CHM surface, which has to be the same as the corresponding KMR example we want to glue (theorem 1.2). To simplify the construction of examples satisfying theorems 1.1 and 1.3, we consider a "no bent" CHM example M_k. In section 3 we perturb $M_k(\xi)$ using the implicit function theorem. We get an infinite dimensional family of minimal surfaces that have three boundaries.



Figure 2. A sketch of a surface in the family of theorem 1.3.

In section 4, we apply an implicit function theorem to solve the Dirichlet problem for the minimal graph equation on a horizontal flat periodic annuli with a disk B removed, prescribing the boundary data on ∂B and the asymptotic direction of the Scherk-type ends. We construct the flat annuli with a disk removed we will glue to the CHM example at its middle planar end. Varying the asymptotic direction of the ends and the flux of the surface, we obtain the pieces of Scherk example we will glue at the top and bottom catenoidal ends of the CHM example (proving theorem 1.1) and to half a KMR example (theorem 1.3).

In section 5, we study the KMR examples $M_{\sigma,\alpha,\beta}$ and describe a conformal parametrization of these examples on a cylinder. We also obtain an expansion of pieces of the KMR examples as the flux of $M_{\sigma,\alpha,\beta}$ becomes horizontal (i.e. near the catenoidal limit). Section 6 is devoted to the study of the mapping properties of the Jacobi operator about such $M_{\sigma,\alpha,\beta}$ near the catenoidal limit. And we apply in section 7 the implicit function theorem to perturb half a KMR example $M_{\sigma,\alpha,0}$ (resp. $M_{\sigma,0,\beta}$), obtaining a family of minimal surfaces asymptotic to half a $M_{\sigma,\alpha,0}$ (resp. $M_{\sigma,0,\beta}$) and whose boundary is a Jordan curve. We prescribe the boundary data of such a surface. We remark that sections 5, 6, 7 are of independent interest. They are devoted to the global analysis on KMR examples.

Finally, we do the end-to-end construction in section 8: we explain how the boundary data of the corresponding minimal surfaces constructed in sections 4, 3 and 7 can be chosen so that the union of these forms smooth minimal surfaces satisfying theorems 1.1, 1.2 and 1.3.

2. A Costa-Hoffman-Meeks type surface with bent catenoidal ends

In this section we recall the result shown in [6] about the existence of a family of minimal surfaces $M_k(\xi)$ close to the Costa-Hoffman-Meeks surface $M_k(0) = M_k$ of genus $k \geqslant 1$, with one planar end and two slightly bent catenoidal ends by an angle ξ.

2.1. **Costa-Hoffman-Meeks surfaces.** We briefly present here the family of CHM surfaces M_k studied in [2, 3, 8, 9, 10]. For each natural $k \geqslant 1$, M_k is a properly embedded minimal surface of genus k and three ends. After suitable rotations and translations, we can assume its ends are horizontal (in particular, they can be ordered by heights) and it enjoys the following properties:

(1) M_k has one middle planar end E_m asymptotic to the plane $\{x_3 = 0\}$, and two catenoidal ends: one top E_t and one bottom E_b, respectively asymptotic to the upper and lower end of a catenoid having as axis of revolution the x_3-axis.

(2) M_k intersects the $\{x_3 = 0\}$ plane in $k+1$ straight lines, which intersect at equal angles $\frac{\pi}{k+1}$ at the origin. The intersection of M_k with any one of the remaining horizontal planes is a single Jordan curve. Thus the intersection of M_k with the upper half-space $\{x_3 > 0\}$ (resp. the lower half-space $\{x_3 < 0\}$) is topologically an open annulus.

(3) The symmetry group of M_k is generated by π-rotations about the $k+1$ lines contained in the surface at height zero, together with the reflection symmetries in vertical planes that bisect those lines. Assume one of such planes of symmetry is the $\{x_2 = 0\}$ plane. In particular, M_k is invariant by the rotation of angle $\frac{2\pi}{k+1}$ about the x_3-axis and by the composition of a rotation by angle $\frac{\pi}{k+1}$ about the x_3-axis with the reflection symmetry across the $\{x_3 = 0\}$ plane.

Now we give a local description of the surfaces M_k near its ends and we introduce coordinates that we will use.

The planar end. The planar end E_m of M_k can be parametrized [6] by

$$X_m(x) = \left(\frac{x}{|x|^2}, u_m(x)\right) \in \mathbb{R}^3, \quad x \in \bar{B}^*_{\rho_0}(0)$$

where $\bar{B}^*_{\rho_0}(0)$ is the punctured closed disk in $\mathbb{R}^2$ of radius $\rho_0 > 0$ small centered at the origin, and $u_m = \mathcal{O}_{C^{2,\alpha}_b}(|x|^{k+1})$ is a solution of

$$|x|^4 \operatorname{div}\left(\frac{\nabla u}{(1+|x|^4\,|\nabla u|^2)^{1/2}}\right) = 0. \tag{1}$$

Moreover, u_m can be extended continuously to the puncture, using Weierstrass representation (in fact, it can be extended as a $C^{2,\alpha}$ function). Here $\mathcal{O}_{C^{n,\alpha}_b}(g)$ denotes a function that, together with its partial derivatives of order less than or equal to $n+\alpha$, is bounded by a constant times g.

If we linearize in $u = 0$ the nonlinear equation (1), we obtain the expression of an operator which is the Jacobi operator about the plane; i.e. $\mathcal{L}_{\mathbb{R}^2} = |x|^4 \Delta_0$. To be more precise, the linearization of (1) gives

$$L_u\, v = |x|^4 \operatorname{div}\left(\frac{\nabla v}{\sqrt{1+|x|^4|\nabla u|^2}} - |x|^4 \nabla u \frac{\nabla u \cdot \nabla v}{\sqrt{(1+|x|^4|\nabla u|^2)^3}}\right). \tag{2}$$

Equation (1) means that the surface Σ_u parametrized by $x \mapsto \left(\frac{x}{|x|^2}, u(x)\right)$ is minimal. We will give the expression of the mean curvature H_{u+v} of Σ_{u+v} in terms of the mean curvature H_u of Σ_u.

Lemma 2.1. *There exists a function Q_u satisfying $Q_u(0,0) = \nabla Q'_u(0,0) = 0$ such that*

$$2H_{u+v} = 2H_u + L_u v + |x|^4 Q_u(\sqrt{|x|^4}\nabla v, \sqrt{|x|^4}\nabla^2 v). \tag{3}$$

Proof. Define $f(t) = \frac{1}{\sqrt{1+|x|^4|\nabla(u+tv)|^2}}$ and apply Taylor expansion. □

Since u satisfies (1), then $H_u = 0$. The minimal surface equation that we will use in the following sections is:

$$|x|^4 \left(\Delta_0 v + \sqrt{1+|x|^4|\nabla u|^2}\left(\bar{L}_u v + Q_u(|x|^2\nabla v, |x|^2\nabla^2 v)\right)\right) = 0, \tag{4}$$

where $\bar{L}_u$ is a second order linear operator whose coefficients are in $\mathcal{O}_{C^{2,\alpha}}\left(|x|^{k+1}\right)$.

The catenoidal ends. We denote by X_c the parametrization of the standard catenoid C whose axis of revolution is the x_3-axis. Its expression is

$$X_c(s,\theta) := (\cosh s\,\cos\theta, \cosh s\,\sin\theta, s) \in \mathbb{R}^3$$

where $(s,\theta) \in \mathbb{R}\times S^1$. The unit normal vector field about C is

$$n_c(s,\theta) := \frac{1}{\cosh s}(\cos\theta, \sin\theta, -\sinh s), \quad (s,\theta)\in\mathbb{R}\times S^1.$$

Up to a dilation, we can assume that the two ends E_t and E_b of M_k are asymptotic to some translated copy of the catenoid parametrized by X_c in the vertical direction. Therefore, E_t and E_b can be parametrized, respectively, by

$$X_t := X_c + w_t\, n_c + \sigma_t\, e_3 \quad \text{in } (s_0,\infty)\times S^1,$$

$$X_b := X_c - w_b\, n_c - \sigma_b\, e_3 \quad \text{in } (-\infty,-s_0)\times S^1,$$

where $\sigma_t, \sigma_b \in \mathbb{R}$, and w_t (resp. w_b) is a function defined in $(s_0,\infty)\times S^1$ (resp. $(-\infty,-s_0)\times S^1$) which tends exponentially fast to 0 as s goes to $+\infty$ (resp. $-\infty$), reflecting the fact that the ends are asymptotic to a catenoidal end.

We recall that the surface parametrized by $X := X_c + w\, n_c$ is minimal if and only if the function w satisfies the minimal surface equation which, for normal graphs over a catenoid has the following form

$$\mathbb{L}_C w + \frac{1}{\cosh^2 s}\left(Q_2\left(\frac{w}{\cosh s}\right) + \cosh s\, Q_3\left(\frac{w}{\cosh s}\right)\right) = 0, \tag{5}$$

where $\mathbb{L}_C$ is the Jacobi operator about the catenoid, i.e.

$$\mathbb{L}_C w = \frac{1}{\cosh^2 s}\left(\partial^2_{ss} w + \partial^2_{\theta\theta} w + \frac{2w}{\cosh^2 s}\right),$$

and Q_2, Q_3 are linear second order differential operators which are bounded in $\mathcal{C}^k(\mathbb{R}\times S^1)$, for every k, and satisfy $Q_2(0) = Q_3(0) = 0$, $\nabla Q_2(0) = \nabla Q_3(0) = 0, \nabla^2 Q_3(0) = 0$ and then:

$$\|Q_j(v_2) - Q_j(v_1)\|_{\mathcal{C}^{0,\alpha}([s,s+1]\times S^1)} \leqslant c\left(\sup_{i=1,2}\|v_i\|_{\mathcal{C}^{2,\alpha}([s,s+1]\times S^1)}\right)^{j-1} \|v_2 - v_1\|_{\mathcal{C}^{2,\alpha}([s,s+1]\times S^1)} \tag{6}$$

for all $s\in\mathbb{R}$ and all v_1, v_2 such that $\|v_i\|_{\mathcal{C}^{2,\alpha}([s,s+1]\times S^1)} \leqslant 1$. The constant $c>0$ does not depend on s.

2.2. **The family of Costa-Hoffman-Meeks surfaces with bent catenoidal ends.** Denote by R_ξ the rotation of angle ξ about the x_2-axis oriented by e_2. Using an elaborate version of the implicit function theorem and following [12] and [17] it is possible to prove the following

Theorem 2.2 ([6]). *There exists $\xi_0 > 0$ and a smooth 1-parameter family of minimal surfaces $\{M_k(\xi) \mid \xi \in (-\xi_0, \xi_0)\}$ such that $M_k(0) = M_k$ and each $M_k(\xi)$ is invariant by the reflection symmetry across the $\{x_2 = 0\}$ plane, has one horizontal planar end E_m and has two catenoidal ends $E_t(\xi), E_b(\xi)$ asymptotic respectively, up to a translation, to the upper and lower end of the catenoid $R_\xi C$ (i.e. the standard catenoid whose axis of revolution is directed by $R_\xi e_3$). Moreover, $E_t(\xi), E_b(\xi)$ can be parametrized respectively by*

$$X_{t,\xi} = R_\xi \left(X_c + w_{t,\xi}\, n_c\right) + \sigma_{t,\xi}\, e_3 + \varsigma_{t,\xi}\, e_1 \tag{7}$$

$$X_{b,\xi} = R_\xi \left(X_c - w_{b,\xi}\, n_c\right) - \sigma_{b,\xi}\, e_3 - \varsigma_{b,\xi}\, e_1 \tag{8}$$

where the functions $w_{t,\xi}, w_{b,\xi}$ and the numbers $\sigma_{t,\xi}, \varsigma_{t,\xi}, \sigma_{b,\xi}, \varsigma_{b,\xi} \in \mathbb{R}$ depend smoothly on ξ and satisfy

$$|\sigma_{t,\xi}-\sigma_t|+|\sigma_{b,\xi}-\sigma_b|+|\varsigma_{t,\xi}|+|\varsigma_{b,\xi}|+\|w_{t,\xi}-w_t\|_{\mathcal{C}^{2,\alpha}_{-2}([s_0,+\infty)\times S^1)}+\|w_{b,\xi}-w_b\|_{\mathcal{C}^{2,\alpha}_{-2}((-\infty,-s_0]\times S^1)} \leqslant c|\xi|,$$

where

$$\|w\|_{\mathcal{C}^{\ell,\alpha}_\delta([s_0,+\infty)\times S^1)} = \sup_{s \geqslant s_0} \left(e^{-\delta s}\, \|w\|_{\mathcal{C}^{\ell,\alpha}([s,s+1]\times S^1)} \right),$$

$$\|w\|_{\mathcal{C}^{\ell,\alpha}_\delta((-\infty,-s_0])\times S^1)} = \sup_{s \leqslant -s_0} \left(e^{-\delta s}\, \|w\|_{\mathcal{C}^{\ell,\alpha}([s-1,s]\times S^1)} \right).$$

For all $s > s_0$ and $\rho < \rho_0$, we define

$$M_k(\xi, s, \rho) := M_k(\xi) - \left(X_{t,\xi}([s,\infty) \times S^1) \cup X_m(B_\rho(0)) \cup X_{b,\xi}((-\infty,-s] \times S^1) \right). \tag{9}$$

The parametrizations of the three ends of $M_k(\xi)$ induce a decomposition of $M_k(\xi)$ into slightly overlapping components: a compact piece $M_k(\xi, s_0+1, \rho_0/2)$ and three noncompact pieces $X_{t,\xi}((s_0,\infty) \times S^1)$, $X_{b,\xi}((-\infty,-s_0) \times S^1)$ and $X_m(\bar{B}_{\rho_0}(0))$.

We define the weighted space of functions on $M_k(\xi)$.

Definition 2.3. *Given $\ell \in \mathbb{N}$, $\alpha \in (0,1)$ and $\delta \in \mathbb{R}$, we define $\mathcal{C}^{\ell,\alpha}_\delta(M_k(\xi))$ as the space of functions in $\mathcal{C}^{\ell,\alpha}_{loc}(M_k(\xi))$ for which the following norm is finite*

$$\begin{aligned}\|w\|_{\mathcal{C}^{\ell,\alpha}_\delta(M_k(\xi))} &:= \|w\|_{\mathcal{C}^{\ell,\alpha}(M_k(\xi,s_0+1,\rho_0/2))} + \|\, w \circ X_m\|_{\mathcal{C}^{\ell,\alpha}(B_{\rho_0}(0))} \\ &+ \|w \circ X_{t,\xi}\|_{\mathcal{C}^{\ell,\alpha}_\delta([s_0,+\infty)\times S^1)} + \|w \circ X_{b,\xi}\|_{\mathcal{C}^{\ell,\alpha}_\delta((-\infty,-s_0]\times S^1)}\end{aligned}$$

and which are invariant by the reflection symmetry across the $\{x_2 = 0\}$ plane, i.e. $w(p) = w(\bar{p})$ for all $p = (p_1, p_2, p_3) \in M_k(\xi)$, where $\bar{p} := (p_1, -p_2, p_3)$.

We remark that there is no weight on the planar end E_m of $M_k(\xi)$. In fact, we can compactify this end and consider a weighted space of functions defined on a two-ended surface. In the next section we will consider normal perturbations of $M_k(\xi)$ by functions $u \in \mathcal{C}^{2,\alpha}_\delta$ and the planar end E_m will be just vertically translated.

The Jacobi operator. The Jacobi operator about $M_k(\xi)$ is

$$\mathbb{L}_{M_k(\xi)} := \Delta_{M_k(\xi)} + |A_{M_k(\xi)}|^2$$

where $|A_{M_k(\xi)}|$ is the norm of the second fundamental form on $M_k(\xi)$.

In the parametrization of the ends of $M_k(\xi)$ introduced above, the volume form $dvol_{M_k(\xi)}$ can be written as $\gamma_t\, ds\, d\theta$ (resp. $\gamma_b\, ds\, d\theta$, $\gamma_m\, dx_1\, dx_2$) near $E_t(\xi)$ (resp. $E_b(\xi)$, E_m). We define globally on $M_k(\xi)$ a smooth function

$$\gamma : M_k(\xi) \longrightarrow [0,\infty)$$

that equals 1 on $M_k(\xi, s_0-1, 2\rho_0)$ and equals γ_t (resp. γ_b, γ_m) on the end $E_t(\xi)$ (resp. $E_b(\xi)$, E_m). Observe that

$$(\gamma \circ X_{t,\xi})(s,\theta) \sim \cosh^2 s \quad \text{on } X_{t,\xi}((s_0,\infty)\times S^1),$$
$$(\gamma \circ X_{b,\xi})(s,\theta) \sim \cosh^2 s \quad \text{on } X_{b,\xi}((-\infty,-s_0)\times S^1),$$
$$(\gamma \circ X_m)(x) \sim |x|^{-4} \quad \text{on } B_{\rho_0}.$$

Granted the above defined spaces, one can check that:

$$\begin{array}{rccc} \mathcal{L}_{\xi,\delta}: & \mathcal{C}^{2,\alpha}_\delta(M_k(\xi)) & \longrightarrow & \mathcal{C}^{0,\alpha}_\delta(M_k(\xi)) \\ & w & \longmapsto & \gamma\, \mathbb{L}_{M_k(\xi)}\,(w) \end{array}$$

is a bounded linear operator. The subscript δ is meant to keep track of the weighted space over which the Jacobi operator is acting. Observe that, the function γ is here to counterbalance the effect of the conformal factor $\frac{1}{\sqrt{|g_{M_k(\xi)}|}}$ in the expression of the Laplacian in the coordinates we use to parametrize the ends of the surface $M_k(\xi)$. This is precisely what is needed to have the operator defined from the space $\mathcal{C}^{2,\alpha}_\delta(M_k(\xi))$ into the target space $\mathcal{C}^{0,\alpha}_\delta(M_k(\xi))$.

To have a better grasp of what is going on, let us linearize the nonlinear equation (5) at $w = 0$. We get the expression of the Jacobi operator about the standard catenoid

$$\mathbb{L}_C := \frac{1}{\cosh^2 s}\left(\partial_s^2 + \partial_\theta^2 + \frac{2}{\cosh^2 s}\right).$$

We can observe that the operator $\cosh^2 s\, \mathbb{L}_C$ maps the space $(\cosh s)^\delta\, \mathcal{C}^{2,\alpha}((s_0,+\infty)\times S^1)$ into the space $(\cosh s)^\delta\, \mathcal{C}^{0,\alpha}((s_0,+\infty)\times S^1)$.

Similarly, if we linearize the nonlinear equation (1) at $u = 0$, we obtain (see (2) with $u = 0$) the expression of the Jacobi operator about the plane

$$\mathbb{L}_{\mathbb{R}^2} := |x|^4\, \Delta_0.$$

Again, the operator $|x|^{-4}\,\mathbb{L}_{\mathbb{R}^2} = \Delta_0$ clearly maps the space $\mathcal{C}^{2,\alpha}(\bar{B}_{\rho_0})$ into the space $\mathcal{C}^{0,\alpha}(\bar{B}_{\rho_0})$. Now, the function γ plays, for the ends of the surface $M_k(\xi)$, the role played by the function $\cosh^2 s$ for the ends of the standard catenoid and the role played by the function $|x|^{-4}$ for the plane. Since the Jacobi operator about $M_k(\xi)$ is asymptotic to $\mathbb{L}_{\mathbb{R}^2}$ at E_m and is asymptotic to $\mathbb{L}_C$ at $E_t(\xi)$ and $E_b(\xi)$, we conclude that the operator $\mathcal{L}_{\xi,\delta}$ maps $\mathcal{C}^{2,\alpha}_\delta(M_k(\xi))$ into $\mathcal{C}^{0,\alpha}_\delta(M_k(\xi))$.

We recall the notion of non degeneracy introduced in [6]:

Definition 2.4. *The surface $M_k(\xi)$ is said to be non degenerate if $\mathcal{L}_{\xi,\delta}$ is injective for all $\delta < -1$.*

It useful to observe that a duality argument in the weighted Lebesgue spaces, implies

$$(\mathcal{L}_{\xi,\delta} \quad \text{is injective}) \quad \Leftrightarrow \quad (\mathcal{L}_{\xi,-\delta} \quad \text{is surjective})$$

provided $\delta \notin \mathbb{Z}$. See [12, 22] for more details.

The non degeneracy of $M_k(\xi)$ follows from the study of the kernel of $\mathcal{L}_{\xi,\delta}$.

The Jacobi fields. It is known that a smooth 1-parameter group of isometries containing the identity generates a Jacobi field, that is a solution of the equation $\mathbb{L}_{M_k(\xi)} u = 0$. The solutions which are invariant under the action of the reflection symmetry across the $\{x_2 = 0\}$ plane, are generated by dilations, vertical translations and horizontal translations along the x_1-axis (we refer [6] for details):

- The group of vertical translations generated by the Killing vector field $\Xi(p) = e_3$ gives rise to the Jacobi field $\Phi^{0,+}(p) := n(p) \cdot e_3$.
- The vector field $\Xi(p) = p$ associated to the 1-parameter group of dilation, generates the Jacobi fields $\Phi^{0,-}(p) := n(p) \cdot p$.
- The Killing vector field $\Xi(p) = e_1$ that generates the group of translations along the x_1-axis is associated to a Jacobi field $\Phi^{1,+}(p) := n(p) \cdot e_1$.
- Finally, we denote by $\Phi^{1,-}(p) := n(p) \cdot (e_2 \times p)$ the Jacobi field associated to the Killing vector field $\Xi(p) = e_2 \times p$ that generates the group of rotations about the x_2-axis.

There are other Jacobi fields we do not take into account because they are not invariant by the reflection symmetry across the $\{x_2 = 0\}$ plane.

With these notations, we define the deficiency space

$$\mathcal{D} := \mathrm{Span}\{\chi_t\, \Phi^{j,\pm}, \chi_b\, \Phi^{j,\pm} \,:\, j = 0, 1\}$$

where χ_t is a cutoff function that is identically equal to 1 on $X_{t,\xi}((s_0 + 1, \infty) \times S^1)$, identically equal to 0 on $M_k(\xi) - X_{t,\xi}((s_0, \infty) \times S^1)$ and that is invariant under the action of the symmetry with respect to the $\{x_2 = 0\}$ plane; and

$$\chi_b(\cdot) := \chi_t(- \cdot).$$

Clearly,

$$\begin{array}{rccc} \tilde{\mathcal{L}}_{\xi,\delta} : & \mathcal{C}^{2,\alpha}_\delta(M_k(\xi)) \oplus \mathcal{D} & \longrightarrow & \mathcal{C}^{0,\alpha}_\delta(M_k(\xi)) \\ & w & \longmapsto & \gamma\, \mathbb{L}_{M_k(\xi)}\,(w) \end{array}$$

is a bounded linear operator, for $\delta < 0$.

A result of S. Nayatani [25, 26] extended by the second author [24] states that any bounded Jacobi field invariant by the reflection symmetry across the $\{x_2 = 0\}$ plane, is linear combination of $\Phi^{0,+}$ and $\Phi^{1,+}$. This fact together with an adaptation to our setting of the linear decomposition lemma proved in [17] for constant mean curvature surfaces (see also [12] for minimal hypersurfaces), allows us to prove the following result.

Proposition 2.5. *We fix $\delta \in (-2, -1)$. Then (reducing ξ_0 if this is necessary) the operator $\tilde{\mathcal{L}}_{\xi,\delta}$, for $|\xi| < \xi_0$, is surjective and has a kernel of dimension* 4.

From that we get the following one about the operator $\mathcal{L}_{\xi,\delta}$

Proposition 2.6. *We fix $\delta \in (1, 2)$. Then (reducing ξ_0 if this is necessary) the operator $\mathcal{L}_{\xi,\delta}$ is surjective and there exists $G_{\xi,\delta}$ a right inverse for $\mathcal{L}_{\xi,\delta}$ that depends smoothly on ξ and in particular whose norm is bounded uniformly as $|\xi| < \xi_0$.*

3. Infinite dimensional family of minimal surfaces close to $M_k(\xi)$

In this section we consider a truncature of $M_k(\xi)$. First we write portions of the ends of $M_k(\xi)$ as vertical graphs over the $\{x_3 = 0\}$ plane.

Lemma 3.1 ([6]). *There exists $\varepsilon_0 > 0$ such that, for all $\varepsilon \in (0, \varepsilon_0)$ and all $|\xi| \leqslant \varepsilon$, an annular part of the ends $E_t(\xi)$, $E_b(\xi)$ and E_m of $M_k(\xi)$ can be written, respectively, as vertical graphs over the annulus $B_{2r_\varepsilon} - B_{r_\varepsilon/2}$ for the functions*

$$\begin{aligned} U_t(r, \theta) &= \sigma_{t,\xi} + \ln(2r) - \xi\, r\, \cos\theta + \mathcal{O}_{\mathcal{C}^\infty_b}(\varepsilon), \\ U_b(r, \theta) &= -\sigma_{b,\xi} - \ln(2r) - \xi\, r\, \cos\theta + \mathcal{O}_{\mathcal{C}^\infty_b}(\varepsilon), \\ U_m(r, \theta) &= \mathcal{O}_{\mathcal{C}^\infty_b}(r^{-(k+1)}). \end{aligned}$$

Here (r, θ) *are the polar coordinates in the* $\{x_3 = 0\}$ *plane. The functions* $\mathcal{O}(\varepsilon)$ *are defined in the annulus* $B_{2\,r_\varepsilon} - B_{r_\varepsilon/2}$ *and are bounded in the* $\mathcal{C}^\infty_b$ *topology by a constant (independent on* ε*) multiplied by* ε*, where the partial derivatives are computed with respect to the vector fields* $r\,\partial_r$ *and* ∂_θ*.*

In particular, a portion of the two catenoidal ends $E_t(\varepsilon/2)$ and $E_b(\varepsilon/2)$ of $M_k(\varepsilon/2)$ are graphs over the annulus $B_{2\,r_\varepsilon} - B_{r_\varepsilon/2} \subset \{x_3 = 0\}$ for functions U_t and U_b. We set

$$s_\varepsilon = -\frac{1}{2}\ln\varepsilon, \quad \rho_\varepsilon = 2\varepsilon^{1/2} \quad \text{and}$$

$$M_k^T(\varepsilon/2) = M_k(\varepsilon/2) - \left(X_{t,\varepsilon/2}((s_\varepsilon, +\infty) \times S^1) \cup X_{b,\varepsilon/2}((-\infty, -s_\varepsilon) \times S^1) \cup X_m(B_{\rho_\varepsilon}(0))\right).$$

We prove, following section 6 in [6], the existence of a family of surfaces close to $M_k^T(\xi)$. In a first step, we modify the parametrization of the ends $E_t(\varepsilon/2), E_b(\varepsilon/2), E_m$, for appropriates values of s, so that, when $r \in [3r_\varepsilon/4, 3r_\varepsilon/2]$ the curves corresponding to the image of

$$\theta \to (r\,\cos\theta, r\,\sin\theta, U_t(r,\theta)), \quad \theta \to (r\,\cos\theta, r\,\sin\theta, U_b(r,\theta)),$$

$$\theta \to (r\,\cos\theta, r\,\sin\theta, U_m(r,\theta))$$

correspond respectively to the curves $\{s = \ln(2r)\}, \{s = -\ln(2r)\}, \{\rho = \frac{1}{r}\}$.

The second step is the modification of the unit normal vector field on $M_k(\varepsilon/2)$ to produce a transverse unit vector field $\tilde{n}_{\varepsilon/2}$ in such a way that it coincides with the normal vector field $n_{\varepsilon/2}$ on $M_k(\varepsilon/2)$, is equal to e_3 on the graph over $B_{3r_\varepsilon/2} - B_{3r_\varepsilon/4}$ of the functions U_t and U_b and interpolate smoothly between the different definitions of $\tilde{n}_{\varepsilon/2}$ in different subsets of $M_k^T(\varepsilon/2)$.

Finally we observe that close to $E_t(\varepsilon/2)$, we can give the following estimate:

$$\left|\cosh^2 s\left(\mathbb{L}_{M_k(\varepsilon/2)} v - \cosh^{-2} s\left(\partial^2_{ss} v + \partial^2_{\theta\theta} v\right)\right)\right| \leqslant c\left|\cosh^{-2} s\, v\right|. \tag{10}$$

This follows easily from (5) together with the fact that $w_{t,\xi}$ (see (7)) decays at least like $\cosh^{-2} s$ on $E_t(\varepsilon/2)$. Similar considerations hold close the bottom end $E_b(\varepsilon/2)$. Near the middle planar end E_m, we observe that the following estimate holds:

$$\left||x|^{-4}\left(\mathbb{L}_{M_k(\varepsilon/2)} v - |x|^4 \Delta_0 v\right)\right| \leqslant c\left||x|^{2k+3}\nabla v\right|. \tag{11}$$

This follows easily from (2) together with the fact that u_m decays at least like $|x|^{k+1}$ on E_m.

The graph of a function u, using the vector field $\tilde{n}_{\varepsilon/2}$, will be a minimal surface if and only if u is a solution of a second order nonlinear elliptic equation of the form

$$\mathbb{L}_{M_k^T(\varepsilon/2)}\, u = \tilde{L}_{\varepsilon/2}\, u + Q_\varepsilon\,(u)$$

where $\mathbb{L}_{M_k^T(\varepsilon/2)}$ is the Jacobi operator about $M_k^T(\varepsilon/2)$, Q_ε is a nonlinear second order differential operator and $\tilde{L}_{\varepsilon/2}$ is a linear operator which takes into account the change of the normal vector field (only for the top and bottom ends) and the change of the parametrization. This operator is supported in a neighborhoods of $\{\pm s_\varepsilon\} \times S^1$, where its coefficients are uniformly bounded by a constant times ε^2, and a neighborhood of $\{\rho_\varepsilon\} \times S^1$ where its coefficients are uniformly bounded by a constant times ε^3.

Now, we consider three even functions $\varphi_t, \varphi_b, \varphi_m \in \mathcal{C}^{2,\alpha}(S^1)$ such that φ_t, φ_b are L^2-orthogonal to 1 and $\theta \to \cos\theta$ while φ_m is L^2-orthogonal to 1. Assume that they satisfy

$$\|\varphi_t\|_{\mathcal{C}^{2,\alpha}} + \|\varphi_b\|_{\mathcal{C}^{2,\alpha}} + \|\varphi_m\|_{\mathcal{C}^{2,\alpha}} \leqslant \kappa\,\varepsilon. \tag{12}$$

We set $\Phi := (\varphi_t, \varphi_b, \varphi_m)$ and we define w_Φ to be the function equal to:

1. $\chi_+(s)\, H_{\varphi_t}(s_\varepsilon - s, \cdot)$ on the image of $X_{t,\varepsilon/2}$, where χ_+ is a cut-off function equal to 0 for $s \leqslant s_0 + 1$ and identically equal to 1 for $s \in [s_0+2, s_\varepsilon]$,
2. $\chi_-(s)\, H_{\varphi_b}(s - s_\varepsilon, \cdot)$ on the image of $X_{b,\varepsilon/2}$, where χ_- is a cut-off function which equals 0 for $s \geqslant -s_0 - 1$ and 1 for $s \in [-s_\varepsilon, -s_0 - 2]$,
3. $\chi_m(r)\, \tilde{H}_{\rho_\varepsilon, \varphi_m}(\cdot, \cdot)$ on the image of X_m, where χ_m is a cut-off function equal to 0 for $r \geqslant \rho_0$ and to 1 for $\rho \in [\rho_\varepsilon, \rho_0/2]$,
4. 0 in the remaining part of the surface $M_k^T(\varepsilon/2)$,

where $\tilde{H}$ and H are, respectively, harmonic extensions of the operators introduced in Propositions 9.2 and 9.4 of Appendix A.

We would like to prove that, under appropriates hypothesis, the graph over $M_k^T(\varepsilon/2)$ of the function $u = w_\Phi + v$ is a minimal surface. This is equivalent to solve the equation:

$$\mathbb{L}_{M_k^T(\varepsilon/2)}(w_\Phi + v) = \tilde{L}_{\varepsilon/2}(w_\Phi + v) + Q_\varepsilon(w_\Phi + v).$$

The resolution of the previous equation is obtained thanks to the following fixed point problem:

$$v = T(\Phi, v) \tag{13}$$

with

$$T(\Phi, v) = G_{\varepsilon/2,\delta} \circ \mathcal{E}_\varepsilon \left(\gamma \left(\tilde{L}_{\varepsilon/2}(w_\Phi + v) - \mathbb{L}_{M_k^T(\varepsilon/2)}\, w_\Phi + Q_\varepsilon(w_\Phi + v) \right) \right),$$

where $\delta \in (1,2)$, the operator $G_{\varepsilon/2,\delta}$ is the right inverse provided in Proposition 2.6 and $\mathcal{E}_\varepsilon$ is a linear extension operator

$$\mathcal{E}_\varepsilon : \mathcal{C}^{0,\alpha}_\delta(M_k^T(\varepsilon/2)) \longrightarrow \mathcal{C}^{0,\alpha}_\delta(M_k(\varepsilon/2)),$$

(here $\mathcal{C}^{0,\alpha}_\delta(M_k^T(\varepsilon/2))$ denotes the space of functions of $\mathcal{C}^{0,\alpha}_\delta(M_k(\varepsilon/2))$ restricted to $M_k^T(\varepsilon/2)$) such that $\mathcal{E}_\varepsilon v = v$ in $M_k^T(\varepsilon/2)$, $\mathcal{E}_\varepsilon v = 0$ in the image of $[s_\varepsilon + 1, +\infty) \times S^1$ by $X_{t,\varepsilon/2}$, in the image of $(-\infty, -s_\varepsilon - 1) \times S^1$ by $X_{b,\varepsilon/2}$ and in the image of $B_{\rho_\varepsilon/2} \times S^1$ by X_m, and $\mathcal{E}_\varepsilon v$ is an interpolation of these values in the remaining part of $M_k(\varepsilon/2)$:

$$(\mathcal{E}_\varepsilon v) \circ X_{t,\varepsilon/2}(s,\theta) = (1 + s_\varepsilon - s)(v \circ X_{t,\varepsilon/2}(s_\varepsilon, \theta)), \quad \text{for} \quad (s,\theta) \in [s_\varepsilon, s_\varepsilon + 1] \times S^1$$

$$(\mathcal{E}_\varepsilon v) \circ X_{b,\varepsilon/2}(s,\theta) = (1 + s_\varepsilon + s)(v \circ X_{b,\varepsilon/2}(s_\varepsilon, \theta)) \quad \text{for} \quad (s,\theta) \in [-s_\varepsilon - 1, -s_\varepsilon] \times S^1$$

$$(\mathcal{E}_\varepsilon v) \circ X_m(\rho,\theta) = \left(\frac{2}{\rho_\varepsilon}\rho - 1 \right)(v \circ X_m(\rho_\varepsilon, \theta)) \quad \text{for} \quad (\rho,\theta) \in [\rho_\varepsilon/2, \rho_\varepsilon] \times S^1.$$

Remark 3.2. *As consequence of the properties of* $\mathcal{E}_\varepsilon$, *if* $\operatorname{supp} v \cap \left(B_{\rho_\varepsilon} - B_{\rho_\varepsilon/2}\right) \neq \emptyset$ *then*

$$\|(\mathcal{E}_\varepsilon v) \circ X_m\|_{\mathcal{C}^{0,\alpha}(\bar{B}_{\rho_0})} \leqslant c\rho_\varepsilon^{-\alpha} \|v \circ X_m\|_{\mathcal{C}^{0,\alpha}(B_{\rho_0} - B_{\rho_\varepsilon})}.$$

This phenomenon of explosion of the norm does not occur near the catenoidal type ends:

$$\|(\mathcal{E}_\varepsilon v) \circ X_{t,\varepsilon/2}\|_{\mathcal{C}^{0,\alpha}([s_0,+\infty)\times S^1)} \leqslant c \|v \circ X_{t,\varepsilon/2}\|_{\mathcal{C}^{0,\alpha}([s_0,s_\varepsilon]\times S^1)}.$$

A similar equation holds for the bottom end.

In the following we will assume $\alpha > 0$ *and close to zero.*

The existence of a solution $v \in \mathcal{C}^{2,\alpha}_\delta(M_k^T(\varepsilon/2))$ for the equation (13) is a consequence of the following result which proves that T is a contracting mapping.

Proposition 3.3. *There exist constants* $c_\kappa > 0$ *and* $\varepsilon_\kappa > 0$*, such that*

$$\|T(\Phi,0)\|_{\mathcal{C}^{2,\alpha}_\delta}(M_k^T(\varepsilon/2)) \leqslant c_\kappa\, \varepsilon^{3/2} \tag{14}$$

and, for all $\varepsilon \in (0,\varepsilon_\kappa)$ *and* $0 < \alpha < \frac{1}{2}$,

$$\|T(\Phi,v_2) - T(\Phi,v_1)\|_{\mathcal{C}^{2,\alpha}_\delta(M_k(\varepsilon/2))} \leqslant \frac{1}{2}\,\|v_2 - v_1\|_{\mathcal{C}^{2,\alpha}_\delta(M_k(\varepsilon/2))},$$

$$\|T(\Phi_2,v) - T(\Phi_1,v)\|_{\mathcal{C}^{2,\alpha}_\delta(M_k(\varepsilon/2))} \leqslant c\varepsilon\,\|\Phi_2 - \Phi_1\|_{\mathcal{C}^{2,\alpha}(S^1)}, \tag{15}$$

where

$$\|\Phi_2 - \Phi_1\|_{\mathcal{C}^{2,\alpha}(S^1)} = \|\varphi_{t,2} - \varphi_{t,1}\|_{\mathcal{C}^{2,\alpha}(S^1)}$$
$$+\|\varphi_{b,2} - \varphi_{b,1}\|_{\mathcal{C}^{2,\alpha}(S^1)} + \|\varphi_{m,2} - \varphi_{m,1}\|_{\mathcal{C}^{2,\alpha}(S^1)}$$

for all $v, v_1, v_2 \in \mathcal{C}^{2,\alpha}_\delta(M_k^T(\varepsilon/2))$ *and satisfying* $\|v\|_{\mathcal{C}^{2,\alpha}_\delta} \leqslant 2\,c_\kappa\,\varepsilon^{3/2}$ *and for all boundary data* $\Phi, \Phi_1, \Phi_2 \in [\mathcal{C}^{2,\alpha}(S^1)]^3$ *satisfying* (12)*.*

Proof. We recall that the Jacobi operator associated to $M_k(\varepsilon/2)$, is asymptotic to the operator of the catenoid near the catenoidal ends, and it is asymptotic to the laplacian near of the planar end. The function w_Φ is identically zero far from the ends where the explicit expression of $\mathbb{L}_{M_k(\varepsilon/2)}$ is not known: this is the reason of our particular choice in the definition of w_Φ. Then from the definition of w_Φ, thanks to Proposition 2.6, to (10) and (11), we obtain the estimate

$$\|\mathcal{E}_\varepsilon\left(\gamma \mathbb{L}_{M_k^T(\varepsilon/2)}\, w_\Phi\right)\|_{\mathcal{C}^{0,\alpha}_\delta(M_k(\varepsilon/2))} =$$
$$\leqslant c\left|\left|\cosh^{-2} s\,(w_\Phi \circ X_{t,\varepsilon/2})\right|\right|_{\mathcal{C}^{0,\alpha}_\delta([s_0+1,s_\varepsilon]\times S^1)} + c\left|\left|\cosh^{-2} s\,(w_\Phi \circ X_{b,\varepsilon/2})\right|\right|_{\mathcal{C}^{0,\alpha}_\delta([-s_\varepsilon,-s_0-1]\times S^1)} +$$
$$c\varepsilon^{-\frac{\alpha}{2}}\left|\left|\rho^{2k+3}\nabla(w_\Phi \circ X_m)\right|\right|_{\mathcal{C}^{0,\alpha}([\rho_\varepsilon,\rho_0]\times S^1)} \leqslant c_\kappa \varepsilon^{3/2}.$$

Using the properties of $\tilde{L}_{\varepsilon/2}$, we obtain

$$\|\mathcal{E}_\varepsilon\left(\gamma \tilde{L}_{\varepsilon/2}\, w_\Phi\right)\|_{\mathcal{C}^{0,\alpha}_\delta(M_k(\varepsilon/2))} \leqslant c\varepsilon\|w_\Phi \circ X_{t,\varepsilon/2}\|_{\mathcal{C}^{0,\alpha}_\delta([s_0+1,s_\varepsilon]\times S^1)}$$
$$+ c\varepsilon\|w_\Phi \circ X_{b,\varepsilon/2}\|_{\mathcal{C}^{0,\alpha}_\delta([-s_\varepsilon,-s_0-1]\times S^1)} + c\varepsilon^{1-\alpha/2}\|w_\Phi \circ X_m\|_{\mathcal{C}^{0,\alpha}([\rho_\varepsilon,\rho_0]\times S^1)} \leqslant c_\kappa \varepsilon^{3/2}.$$

As for the last term, we recall that the operator Q_ε has two different expressions if we consider the catenoidal type end and the planar end (see (5) and (4)). It holds that

$$\|\mathcal{E}_\varepsilon\left(\gamma Q_\varepsilon\left(w_\Phi\right)\right)\|_{\mathcal{C}^{0,\alpha}_\delta(M_k(\varepsilon/2))} \leqslant c_k \varepsilon^{3/2}.$$

Details are left to the reader. □

Theorem 3.4. *Let be* $B := \{w \in \mathcal{C}^{2,\alpha}_\delta(M_{k,\varepsilon}) \,|\, ||w||_{2,\alpha} \leqslant 2c_\kappa\varepsilon^{3/2}\}$*. Then the nonlinear mapping* T *defined above has a unique fixed point* v *in* B*.*

Proof. The previous lemma shows that, if ε is chosen small enough, the nonlinear mapping T is a contraction mapping from the ball B of radius $2c_\kappa\varepsilon^{3/2}$ in $\mathcal{C}^{2,\alpha}_\delta(M_k^T(\varepsilon/2))$ into itself. This value follows from the estimate of the norm of $T(\Phi,0)$. Consequently thanks to Schäuder fixed point theorem, T has a unique fixed point w in this ball. □

This argument provides a minimal surface $M_k^T(\varepsilon/2,\Phi)$ which is close to $M_k^T(\varepsilon/2)$ and has three boundaries. This surface is, close to its upper and lower boundary, a vertical graph over the annulus $B_{r_\varepsilon} - B_{r_{\varepsilon/2}}$ whose parametrization is, respectively, given by

$$U_t(r,\theta) = \sigma_{t,\varepsilon/2} + \ln(2r) + \frac{\varepsilon}{2} r\cos\theta + H_{\varphi_t}(s_\varepsilon - \ln 2r, \theta) + V_t(r,\theta),$$

$$U_b(r,\theta) = -\sigma_{b,\varepsilon/2} - \ln(2r) + \frac{\varepsilon}{2} r\cos\theta + H_{\varphi_b}(s_\varepsilon - \ln 2r, \theta) + V_b(r,\theta),$$

where $s_\varepsilon = -\frac{1}{2}\ln\varepsilon$. The boundaries of the surface correspond to $r_\varepsilon = \frac{1}{2}\,\varepsilon^{-1/2}$. Nearby the middle boundary the surface is a vertical graph over the annulus $B_{r_\varepsilon} - B_{r_\varepsilon/2}$, Its parametrization is

$$U_m(r,\theta) = \tilde{H}_{\rho_\varepsilon,\varphi_m}(1/r,\theta) + V_m(r,\theta),$$

where $\rho_\varepsilon = 2\varepsilon^{1/2}$. All the functions V_i for $i = t, b, m$ depend non linearly on ε, φ.

Lemma 3.5. *The functions* $V_i(\varepsilon,\varphi_i)$, *for* $i = t, b$, *satisfy* $\|V_i(\varepsilon,\varphi_i)(r_\varepsilon\cdot,\cdot)\|_{\mathcal{C}^{2,\alpha}(\bar{B}_2 - B_{1/2})} \leqslant c\varepsilon$ *and*

$$\|V_i(\varepsilon,\varphi_{i,2})(r_\varepsilon\cdot,\cdot) - V_i(\varepsilon,\varphi_{i,1})(r_\varepsilon\cdot,\cdot)\|_{\mathcal{C}^{2,\alpha}(\bar{B}_2 - B_{1/2})} \leqslant c\varepsilon^{1-\delta/2}\|\varphi_{i,2} - \varphi_{i,1}\|_{\mathcal{C}^{2,\alpha}} \tag{16}$$

The function $V_m(\varepsilon,\varphi)$ *satisfies* $\|V_m(\varepsilon,\varphi)(\rho_\varepsilon\cdot,\cdot)\|_{\mathcal{C}^{2,\alpha}(\bar{B}_2 - B_{1/2})} \leqslant c\varepsilon$ *and*

$$\|V_m(\varepsilon,\varphi_{m,2})(\rho_\varepsilon\cdot,\cdot) - V_m(\varepsilon,\varphi_{m,1})(\rho_\varepsilon\cdot,\cdot)\|_{\mathcal{C}^{2,\alpha}(\bar{B}_2 - B_{1/2})} \leqslant c\varepsilon\|\varphi_{m,2} - \varphi_{m,1}\|_{\mathcal{C}^{2,\alpha}} \tag{17}$$

Proof. The wanted estimates follow from

$$\|V_i(\varepsilon,\varphi_2)(\cdot,\cdot) - V_i(\varepsilon,\varphi_1)(\cdot,\cdot)\|_{\mathcal{C}^{2,\alpha}(\bar{B}_{2r_\varepsilon} - B_{r_\varepsilon/2})} \leqslant ce^{\delta s_\varepsilon}\|T(\Phi_2,V_i) - T(\Phi_1,V_i)\|_{\mathcal{C}^{2,\alpha}_\delta(E_i(\varepsilon/2))},$$

for $i = t, b$, together with

$$\|V_m(\varepsilon,\varphi_2)(\cdot,\cdot) - V_m(\varepsilon,\varphi_1)(\cdot,\cdot)\|_{\mathcal{C}^{2,\alpha}(\bar{B}_{2\rho_\varepsilon} - B_{\rho_\varepsilon/2})} \leqslant c\|T(\Phi_2,V_m) - T(\Phi_1,V_m)\|_{\mathcal{C}^{2,\alpha}(E_m)}$$

and the estimate (15) of Proposition 3.3. □

4. An infinite family of Scherk type minimal surfaces close to a horizontal periodic flat annuli

This section has two purposes. The first one is to find an infinite family of minimal surfaces close to a horizontal periodic flat annuli Σ with a disk D_s removed. The surfaces of this family have two horizontal Scherk-type ends E_1, E_2 and will be glued on the middle planar end of a CHM surface. We will prescribe the boundary data φ on ∂D_s. Assume the period T of Σ points to the x_2-direction. Then E_1, E_2 have the x_1-axis as asymptotic direction.

The second and more general purpose of this section is to show the existence of an infinite family of minimal graphs over $\Sigma - D_s$ whose ends have slightly modified asymptotic directions. When the asymptotic directions are not horizontal, these surfaces are close to half a Scherk surface, seen as a graph over $\Sigma - D_s$ (see Figure 2, above). A piece of one such surface will be glued to the catenoidal ends of a CHM example M_k and to an end of a KMR example $M_{\sigma,0,0}$. We will prescribe the boundary data on ∂D_s. Since we need to prescribe the flux along ∂D_s, we will modify the asymptotic direction of the ends and we will choose $|T|$ large.

4.1. Scherk type ends.

We parametrize conformally the annulus $\Sigma \subset \mathbb{R}^3/T$ on $\mathbb{C}^*$, with the notation $(x_1, x_2, x_3) = (x_1 + ix_2, x_3)$, by the mapping

$$A(w) = \left(-\frac{|T|}{2\pi}\ln(w), 0\right), \quad w \in \mathbb{C}^*.$$

The horizontal Scherk-type end E_1 described above can be written as the graph of a function $h_1 \in \mathcal{C}^{2,\alpha}(B_r^*(0))$, where $B_r^*(0)$ is the punctured disk $B_r(0) - \{0\}$ of radius $r \in (0,1)$ centered at the origin. The function $h_1(w)$ is bounded and extends to the puncture (see [7]). The end E_1 can be parameterized by

$$X_1(w) = A(w) + h_1(w)e_3 = \left(-\frac{|T|}{2\pi}\ln(w), h_1(w)\right) \in \mathbb{R}^3/T, \quad w \in B_r^*(0),$$

in the orthonormal frame $\mathcal{F} = (e_1, e_2, e_3)$. The asymptotic direction of the end is e_1.

The horizontal Scherk-type end E_2 can be similarly parameterized in $\mathbb{C} - B_{r^{-1}}(0)$. Via an inversion, we can parametrize E_2 by

$$X_2(w) = \left(-\frac{|T|}{2\pi}\ln(w), h_2(w)\right) \in \mathbb{R}^3/T, \quad w \in B_r^*(0),$$

in the frame $\mathcal{F}^- = (-e_1, -e_2, e_3)$, where $h_2 \in \mathcal{C}^{2,\alpha}(B_r^*(0))$ is a bounded function which can be extended to the puncture. Now the asymptotic direction of the end is $-e_1$.

Let us now parametrize a general Scherk-type end, non necessarily horizontal. Let R_θ denote a rotation in $\mathbb{R}^3/T$ by angle θ about the x_2-axis (oriented by e_2). We can paremetrize a non necessarily horizontal Scherk-type end $\widetilde{E}_1$ with asymptotic direction $\cos\theta_1\, e_1 + \sin\theta_1\, e_3$ and limit normal vector $R_{\theta_1}(e_3)$, with $\theta_1 \in [0, \pi/2)$, by

$$\widetilde{X}_1(z) = \left(-\frac{|T|}{2\pi}\ln(z), \widetilde{h}_1(z)\right), z \in B_r^*(0)$$

in the frame $\mathcal{F}(\theta_1) = R_{\theta_1}\mathcal{F}$, where $\widetilde{h}_1 \in \mathcal{C}^{2,\alpha}(B_r^*(0))$ is a bounded function which can be extended to the origin.

Finally, a Scherk-type end $\widetilde{E}_2$ with asymptotic direction $-\cos\theta_2\, e_1 + \sin\theta_2\, e_3$ and limit normal vector $R_{-\theta_2}(e_3)$, with $\theta_2 \in [0, \pi/2)$, can be parameterized by

$$\widetilde{X}_2(z) = \left(-\frac{|T|}{2\pi}\ln(z), \widetilde{h}_2(z)\right), z \in B_r^*(0)$$

in the frame $\mathcal{F}^-(\theta_2) = R_{-\theta_2}\mathcal{F}^-$, where $\widetilde{h}_2 \in \mathcal{C}^{2,\alpha}(B_r^*(0))$ is a bounded function which can be extended to the origin.

4.2. **Construction of the infinite families.** Given $r \in (0,1)$ and $\Theta = (\theta_1, \theta_2) \in [0, \theta_0]^2$, with $\theta_0 > 0$ small, we denote by $A_\Theta : \mathbb{C}^* \to \mathbb{R}^3/T$ the immersion obtained as smooth interpolation of

$$A_\Theta(z) = \begin{cases} (R_{\theta_1} \circ A)(z) & \text{if } |z| < r/2, \\ A(z) & \text{if } r < |z| < r^{-1}, \\ (R_{-\theta_2} \circ A)(z) & \text{if } |z| > 2r^{-1}. \end{cases}$$

Let N_Θ be the vector field obtained as smooth interpolation of $R_{\theta_1}(e_3)$ on $\{|z| < r/2\}$, e_3 on $\{r < |z| < r^{-1}\}$ and $R_{-\theta_2}(e_3)$ on $\{|z| > 2r^{-1}\}$. For any $h \in C^{2,\alpha}(\bar{\mathbb{C}})$, we define the immersion

$$X_{\Theta,h}(z) = A_\Theta(z) + h(z)N_\Theta(z), \quad z \in \mathbb{C}^*.$$

The immersion $X_{\Theta,h}$ has two Scherk-type ends E_1, E_2 with asymptotic directions $\cos\theta_1\, e_1 + \sin\theta_1\, e_3$ and $-\cos\theta_2\, e_1 + \sin\theta_2\, e_3$, respectively.

At the end E_1 (resp. E_2), $X_{\Theta,h}(z) = A(z) + h_1(z)e_3$ in the orthogonal frame $\mathcal{F}(\theta_1)$ (resp. $X_{\Theta,h}(z) = A(z^{-1}) + h_2(z)e_3$ in the frame $\mathcal{F}^-(\theta_2)$), with $z \in B_r^*(0)$, where $h_1(z) = h(z)$ and $h_2(z) = h(z^{-1})$. In [7] has been proven that the mean curvature of $X_{\Theta,h}$ at E_i is given in terms of the z-coordinate by

$$2H = \frac{4\pi^2|z|^2}{|T|^2}\mathrm{div}_0(P^{-1/2}\nabla_0 h_i),$$

where $P = 1 + \frac{4\pi^2|z|^2}{|T|^2}\|\nabla_0 h_i\|_0^2$ and the subindex $\cdot_0$ means that the corresponding object is computed with respect to the flat metric of the z-plane. We denote by λ the smooth function without zeroes defined by $\lambda(z) = \frac{|T|^2}{4\pi^2|z|^2}$, for $z \in B_r^*(0)$. Then at E_i we have

$$2\lambda H = P^{-1/2}\Delta_0 h_i - \frac{1}{2}P^{-3/2}\langle\nabla_0 P, \nabla_0 h_i\rangle_0.$$

So the mean curvature at the end E_i vanishes if h_i satisfies the equation

$$\Delta_0 h - \frac{1}{2P}\langle \nabla_0 P, \nabla_0 h\rangle_0 = 0. \tag{18}$$

Definition 4.1. *Given $k \in \mathbb{N}$ and $\alpha \in (0,1)$ we define $C^{k,\alpha}(\bar{\mathbb{C}})$ as the space of functions $u \in C^{k,\alpha}_{loc}(\bar{\mathbb{C}})$ such that*

$$||u||_{C^{k,\alpha}(\bar{\mathbb{C}})} := [u]_{k,\alpha,\bar{\mathbb{C}}} < +\infty,$$

where $[u]_{k,\alpha,\bar{\mathbb{C}}}$ denotes the usual $C^{k,\alpha}$ Hölder norm on $\bar{\mathbb{C}}$.

Let B_s be the disk of radius s in $\mathbb{C}^*$ such that $D_s = A(B_s) \subset \Sigma = \{z \in \mathbb{C} \,|\, -|T| < 2y \leqslant |T|\}$ is a geodesic disk centered at the origin of $\mathbb{R}^3/T$. Denote by $C^{k,\alpha}(\bar{\mathbb{C}} - B_s)$ the space of functions in $C^{k,\alpha}(\bar{\mathbb{C}})$ restricted to $\bar{\mathbb{C}} - B_s$. We denote by $H(\Theta, h)$ the mean curvature of $X_{\Theta,h}$, and $\bar{H}(\Theta, h) = H(\Theta, h)$, where λ is the smooth function defined in a neighbourhood at each puncture by $\lambda(z) = \frac{|T|^2}{4\pi^2|z|^2}$. Lemma 4.1 of [7] shows that

$$\bar{H} : \mathbb{R}^2 \times \mathcal{C}^{2,\alpha}(\bar{\mathbb{C}} - B_s) \longrightarrow \mathcal{C}^{0,\alpha}(\bar{\mathbb{C}} - B_s)$$

is an analytical operator. Denote by $\mathcal{L}_\Theta$ the Jacobi operator about A_Θ. We set $\bar{\mathcal{L}}_\Theta = \lambda \mathcal{L}_\Theta$.

Remark 4.2. *The operators H and $\mathcal{L}_\Theta$ are the mean curvature operator and the Jacobi operator with respect to the metric $|dz|^2$ of $\bar{\mathbb{C}}$. Defining the operators $\bar{H} = \lambda H$ and $\bar{\mathcal{L}}_\Theta = \lambda \mathcal{L}_\Theta$ means to consider a different metric on $\bar{\mathbb{C}}$. Actually $\bar{H}$ and $\bar{\mathcal{L}}_\Theta$ are the mean curvature operator and Jacobi operator with respect to the metric $g_\lambda = |dz|^2/\lambda$. From the definition of λ, it follows that the volume of $\bar{\mathbb{C}}$ with respect this metric is finite.*

$\bar{\mathcal{L}}_\Theta$ is a second order linear elliptic operator satisfying $|\bar{\mathcal{L}}_\Theta u - \Delta u| \leqslant c(|\theta_1| + |\theta_2|)|u|$ and the coefficients of $F_\Theta = \Delta - \bar{\mathcal{L}}_\Theta$ have compact support.

Now we fix $s_0 > 0$. Given $\varepsilon > 0$ and $|T| \in [4/\sqrt{\varepsilon}, +\infty)$ large enough, we choose $s \in (0, s_0)$ such that $D_s = A(B_s)$ is the geodesic disk of radius $1/2\sqrt{\varepsilon}$ centered at the origin.

Proposition 4.3. *There exists $\varepsilon_0 > 0$ and $\eta_0 > 0$ such that for every $\varepsilon \in (0, \varepsilon_0)$ and every $|T| \in (\eta_0, +\infty)$, there exists an operator*

$$G_{\varepsilon,|T|} : \mathcal{C}^{0,\alpha}(\bar{\mathbb{C}} - B_s) \longrightarrow \mathcal{C}^{2,\alpha}(\bar{\mathbb{C}} - B_s)$$

such that, given $f \in \mathcal{C}^{0,\alpha}(\bar{\mathbb{C}} - B_s)$, $w = G_{\varepsilon,|T|} f$ satisfies

$$\begin{cases} \Delta w = f & \text{on} \quad \bar{\mathbb{C}} - B_s \\ w \in \text{Span}\{1\} & \text{on} \quad \partial B_s \end{cases}$$

and $||w||_{\mathcal{C}^{2,\alpha}} \leq c||f||_{\mathcal{C}^{0,\alpha}}$ for some constant $c > 0$ which does not depend on $\varepsilon, |T|$.

Proof. Let be u a solution of $\Delta u = f$ on $\bar{\mathbb{C}} - B_s$ with $u = 0$ on ∂B_s. We set $w = u - \int_{\bar{\mathbb{C}} - B_s} u \, dvol_{g_\lambda}$. We recall that the metric in use on $\bar{\mathbb{C}}$ is given by $g_\lambda = |dz|^2/\lambda$. With respect to this metric $vol(\bar{\mathbb{C}} - B_s) < +\infty$ and $\int_{\bar{\mathbb{C}} - B_s} u \, dvol_{g_\lambda} < \infty$. So the function w is well defined and then $\int_{\bar{\mathbb{C}} - B_s} w \, dvol_{g_\lambda} = 0$ and $w \in \text{Span}\{1\}$ on ∂B_s. If the theorem is false, there is a sequence of functions f_n, of solutions w_n and of real numbers s_n such that

$$\sup_{\bar{\mathbb{C}} - B_{s_n}} |f_n| = 1, \quad A_n := \sup_{\bar{\mathbb{C}} - B_{s_n}} |w_n| \to +\infty \text{ as } n \to +\infty,$$

where $s_n \in [0, s_0]$. Now we set $\tilde{w}_n := w_n / A_n$. Elliptic estimates imply that s_n and $\tilde{w}_n$ converge up to a subsequence, respectively, to $s_\infty \in [0, s_0]$ and to $\tilde{w}_\infty$ on $\bar{\mathbb{C}} - B_{s_\infty}$. This function satisfies

$$\Delta \tilde{w}_\infty = 0.$$

Then $\tilde{w}_\infty = \text{const}$ on $\bar{\mathbb{C}} - B_{s_\infty}$ and $\int_{\bar{\mathbb{C}}-B_{s_\infty}} \tilde{w}_\infty \, dvol_{g_\lambda} = 0$, a contradiction with $\sup \widetilde{|w_\infty|} = 1$. □

Now we fix $|T| \geqslant 4/\sqrt{\varepsilon}$, $\Theta \in (0, \varepsilon)^2$, $s_\varepsilon = \frac{1}{2\sqrt{\varepsilon}}$ and let $\varphi \in \mathcal{C}^{2,\alpha}(S^1)$ be even (or odd) L^2-orthogonal to the function $z \to 1$, with $\|\varphi\|_{\mathcal{C}^{2,\alpha}(S^1)} \leqslant \kappa\varepsilon$. Let $w_\varphi = \tilde{H}_{s_\varepsilon,\varphi}$ (see proposition 9.2) be the unique bounded harmonic extension of φ. We would like to solve the minimal surface equation $H(\Theta, v + w_\varphi) = 0$ with fixed boundary data φ, prescribed asymptotic direction Θ and with period $|T|$. Then we have to solve the equation:

$$\Delta v = F_\Theta(v + w_\varphi) + Q_\Theta(v + w_\varphi)$$

with Q_Θ a quadratic term such that $|Q_\Theta(v_1) - Q_\Theta(v_2)| \leqslant c|v_1 - v_2|^2$. The resolution of the previous equation is obtained by showing the existence of a fixed point

$$v = S(\Theta, \varphi, v)$$

where

$$S(\Theta, \varphi, v) = G_{\varepsilon,|T|}(F_\Theta(v + w_\varphi) + Q_\Theta(v + w_\varphi)).$$

Proposition 4.4. *There exists $c_\kappa > 0$ and $\varepsilon_\kappa > 0$ such that for all $|T| \geqslant 4/\sqrt{\varepsilon}$ we have*

$$||S(\Theta, \varphi, 0)||_{\mathcal{C}^{2,\alpha}} \leqslant c_\kappa \varepsilon^2$$

and for all $\varepsilon \in (0, \varepsilon_\kappa)$,

$$||S(\Theta, \varphi, v_1) - S(\Theta, \varphi, v_2)||_{\mathcal{C}^{2,\alpha}} \leqslant \frac{1}{2}||v_2 - v_1||_{\mathcal{C}^{2,\alpha}}$$

$$||S(\Theta, \varphi_1, v) - S(\Theta, \varphi_2, v)||_{\mathcal{C}^{2,\alpha}} \leqslant c\varepsilon||\varphi_2 - \varphi_1||_{\mathcal{C}^{2,\alpha}}$$

for all $v, v_1, v_2 \in \mathcal{C}^{2,\alpha}(\bar{\mathbb{C}} - B_s)$ and satisfying $||v||_{\mathcal{C}^{2,\alpha}} \leqslant 2c_\kappa\varepsilon^2$, for all boundary data $\varphi, \varphi_1, \varphi_2 \perp 1$, whose norm is bounded by a constant κ times ε and for all $\Theta = (\theta_1, \theta_2)$ such that $|\theta_1| + |\theta_2| \leqslant \varepsilon$.

Proof. Using the proposition 4.3, the inequality $|\bar{\mathcal{L}}u - \Delta u| \leqslant c(|\theta_1| + |\theta_2|)|u|$ and the quadratic behavior of Q_Θ we derive the estimate of the proposition. The details of the proof are left to the reader. □

On the set $B_{2s_\varepsilon} - B_{s_\varepsilon}$, the function $U = v + \tilde{H}_{s_\varepsilon,\varphi}$ is the solution of the equation (18). Using the vertical translation $c_0 e_3$ we can fix the value $c_0 + \varphi$ at the boundary:

$$U = c_0 + \tilde{H}_{s_\varepsilon,\varphi} + v.$$

Using Schäuder estimate for the equation on a fixed bounded domain

$$||v(\varphi_1) - v(\varphi_2)||_{\mathcal{C}^{2,\alpha}(\bar{\mathbb{C}} - B_s)} \leqslant c_\kappa \varepsilon ||\varphi_1 - \varphi_2||_{\mathcal{C}^{2,\alpha}(S^1)}.$$

This can be done uniformly in (θ_1, θ_2). We observe that the function U grows logarithmically close $\partial B_{s_\varepsilon}$. The hypothesis of orthogonality of φ to the constant function, implies that the function w_φ enjoys the same property and is bounded. It is not the case of v which can be seen as the sum of a bounded function which is orthogonal to the constant and a function of the form $c \ln(r/s_\varepsilon)$, where $c = c(|T|, \theta_1, \theta_2)$, defined in a neighbourhood of $\partial B_{s_\varepsilon}$. We are able to determine c using flux formula. Let γ_1, γ_2 be two closed curves in $\bar{\Sigma}/T$ chosen in such a way to correspond by the conformal mapping to the boundaries of two circular neighbourhoods N_1, N_2 of the punctures corresponding to the ends with linear growth. Let $\mathcal{S} = \bar{\mathbb{C}} - (B_{s_\varepsilon} \cup N_1 \cup N_2)$. Now we observe that, being X the parametrization of a minimal surface, then the following equality holds:

$$\int_{\mathcal{S}} \Delta X = 0.$$

Thanks to the divergence theorem, if $\Gamma = \partial \mathcal{S}$, then

$$\int_{\mathcal{S}} \Delta X = \int_{\Gamma} \frac{\partial X}{\partial \eta} ds = \int_{\gamma_1} \frac{\partial X}{\partial \eta} ds + \int_{\gamma_2} \frac{\partial X}{\partial \eta} ds + \int_{\partial B_{s_\varepsilon}} \frac{\partial X}{\partial \eta} ds = 0,$$

where η denotes the conormal along Γ. This equality implies

$$\int_{\partial B_{s_\varepsilon}} \frac{\partial U}{\partial \eta} ds = -\sin\theta_1 |T| - \sin\theta_2 |T|.$$

By integration we can conclude that on $B_{2s_\varepsilon} - B_{s_\varepsilon}$, it holds that

$$U = -\frac{|T|}{2\pi}(\sin\theta_1 + \sin\theta_2)\ln(r/s_\varepsilon) + c_0 + w_\varphi + v^\perp.$$

with $v^\perp \perp 1$. Now we choose $|T|$ such that $\frac{|T|}{2\pi}(\sin\theta_1 + \sin\theta_2) = 1$.

We observe that, if $\theta_2 = \theta_1 = 0$, we can state that there exists an infinite family of minimal surfaces which are close to the surface $\Sigma - D_{s_\varepsilon}$. We denote by $S_m(\varphi)$ one of such surfaces. It can be seen as the graph about $B_{2s_\varepsilon} - B_{s_\varepsilon}$ of the function

$$\bar{U}_m(r,\theta) = c_0 + \tilde{H}_{s_\varepsilon,\varphi}(r,\theta) + \bar{V}_m$$

where $V_m = \mathcal{O}_{C_b^{2,\alpha}}(\varepsilon)$ and it satisfies

$$||\bar{V}_m(\varphi_1) - \bar{V}_m(\varphi_2)||_{\mathcal{C}^{2,\alpha}(B_{2s_\varepsilon} - B_{s_\varepsilon})} \leqslant c_\kappa \varepsilon ||\varphi_1 - \varphi_2||_{\mathcal{C}^{2,\alpha}(S^1)}, \tag{19}$$

for $\varphi_2, \varphi_1 \in C^{2,\alpha}(S^1)$.

If $(\theta_2, \theta_1) \neq 0$, we can state that there exists an infinite family of minimal surfaces which are close to the periodic Scherk type example. After a vertical translation, they can be seen as the graph about $B_{2s_\varepsilon} - B_{s_\varepsilon}$ of the function

$$\bar{U}_t(r,\theta) = -\ln(2r) + c_0 + \tilde{H}_{s_\varepsilon,\varphi}(r,\theta) + \bar{V}_t$$

where $\bar{V}_t = \mathcal{O}_{C_b^{2,\alpha}}(\varepsilon)$ and it satisfies

$$||\bar{V}_t(\varphi_1) - \bar{V}_t(\varphi_2)||_{\mathcal{C}^{2,\alpha}(B_{2s_\varepsilon} - B_{s_\varepsilon})} \leqslant c_\kappa \varepsilon ||\varphi_1 - \varphi_2||_{\mathcal{C}^{2,\alpha}(S^1)}, \tag{20}$$

for $\varphi_2, \varphi_1 \in C^{2,\alpha}(S^1)$.

Remark 4.5. *If the boundary data φ is an even function, it is clear the surfaces we have just described are symmetric with respect to the vertical plane $x_2 = 0$. Instead, if the boundary data φ is an odd function and $\theta_1 = \theta_2$ the surfaces are symmetric with respect to the vertical plane $x_1 = 0$.*

5. KMR EXAMPLES

In this section, we briefly present the *KMR examples* $M_{\sigma,\alpha,\beta}$ studied in [13, 14, 21, 30] (also called *toroidal halfplane layers*), which are the only properly embedded doubly periodic minimal surfaces with genus one and finitely many parallel (Scherk-type) ends in the quotient (see [27]).

For each $\sigma \in (0, \frac{\pi}{2})$, $\alpha \in [0, \frac{\pi}{2}]$ and $\beta \in [0, \frac{\pi}{2}]$ with $(\alpha, \beta) \neq (0, \sigma)$, consider the rectangular torus $\Sigma_\sigma = \left\{(z,w) \in \overline{\mathbb{C}}^2 \mid w^2 = (z^2 + \lambda^2)(z^2 + \lambda^{-2})\right\}$, where $\lambda = \lambda(\sigma) = \cot\frac{\sigma}{2} > 1$. The KMR example $M_{\sigma,\alpha,\beta}$ is determined by its Gauss map g and the differential of its height function h, which are defined on Σ_σ and given by:

$$g(z,w) = \frac{az+b}{i(\overline{a}-\overline{b}z)}, \qquad dh = \mu\,\frac{dz}{w},$$

with

$$\begin{aligned} a = a(\alpha,\beta) &= \cos\tfrac{\alpha+\beta}{2} + i\cos\tfrac{\alpha-\beta}{2}; \\ b = b(\alpha,\beta) &= \sin\tfrac{\alpha-\beta}{2} + i\sin\tfrac{\alpha+\beta}{2}; \qquad (21) \\ \mu = \mu(\sigma) &= \tfrac{\pi\csc\sigma}{\mathcal{K}(\sin^2\sigma)}, \end{aligned}$$

where $\mathcal{K}(m) = \int_0^{\frac{\pi}{2}} \frac{1}{\sqrt{1-m\sin^2 u}}\,du$, $0<m<1$, is the complete elliptic integral of first kind. Such μ has been chosen so that the vertical part of the flux of $M_{\sigma,\alpha,\beta}$ along any horizontal level section equals 2π.

Remark 5.1. *The following statements give us a better understanding of the geometrical meaning of a, b defined above:*

- (i) $b \to 0$ *if and only if* $\alpha \to 0$ *and* $\beta \to 0$*, in which case* $a \to 1+i$.
- (ii) $|b|^2 + |a|^2 = 2$.
- (iii) *When* $\beta = 0$*, then* $a = (1+i)\cos\frac{\alpha}{2}$ *and* $b = (1+i)\sin\frac{\alpha}{2}$*. In particular,* $b = \mathcal{O}(\alpha)$.
- (iv) *When* $\alpha = 0$*, then* $a = (1+i)\cos\frac{\beta}{2}$ *and* $b = (-1+i)\sin\frac{\beta}{2}$*. In particular,* $b = \mathcal{O}(\beta)$.
- (v) *In general,* $\left|\frac{b}{a}\right| = \tan\frac{\varphi}{2}$*, where* φ *is the angle between the North Pole* $(0,0,1) \in \mathbb{S}^2$ *and the pole of* g *seen in* $\mathbb{S}^2$ *via the inverse of the stereographic projection.*

The KMR example $M_{\sigma,\alpha,\beta}$ can be parametrized on Σ_σ by the immersion $X = (X_1, X_2, X_3) = \Re \int \mathcal{W}$, where $\mathcal{W}$ is the Weierstrass form:

$$\mathcal{W} = \left(\frac{1}{2}\left(\frac{1}{g} - g\right) dh, \frac{i}{2}\left(\frac{1}{g} + g\right) dh, dh \right).$$

The ends of $M_{\sigma,\alpha,\beta}$ correspond to the punctures $\{A, A', A'', A'''\} = g^{-1}(\{0,\infty\})$, and the branch values of g are those with $w = 0$, i.e.

$$D = (-i\lambda, 0),\ D' = (i\lambda, 0),\ D'' = (\tfrac{i}{\lambda}, 0),\ D''' = (-\tfrac{i}{\lambda}, 0). \qquad (22)$$

Seen in $\mathbb{S}^2$, these points form two pairs of antipodal points: $D'' = -D$ and $D''' = -D'$ and each KMR example can be given in terms of the branch values of its Gauss map (see [30]).

In [30], it is proven that the above Weierstrass data define a properly embedded minimal surface $M_{\sigma,\alpha,\beta}$ invariant by two independent translations: the translation by the period T at its ends, and the translation by the period $\widetilde{T}$ along a homology class. Moreover, the group of isometries $\mathrm{Iso}(M_{\sigma,\alpha,\beta})$ of $M_{\sigma,\alpha,\beta}$ always contains a subgroup isomorphic to $(\mathbb{Z}/2\mathbb{Z})^2$, with generators $\mathcal{D}$ (corresponding to the deck transformation $(z,w) \mapsto (z,-w)$), which represents in $\mathbb{R}^3$ a central symmetry about any of the four branch points of the Gauss map of $M_{\sigma,\alpha,\beta}$; and $\mathcal{F}$, which consists of a translation by $\frac{1}{2}(T + \widetilde{T})$. In particular, the ends of $M_{\sigma,\alpha,\beta}$ are equally spaced.

We are going to focus on two more symmetric subfamilies of KMR examples: $\{M_{\sigma,\alpha,0} \mid 0 < \sigma < \frac{\pi}{2},\ 0 \leqslant \alpha \leqslant \frac{\pi}{2}\}$ and $\{M_{\sigma,0,\beta} \mid 0 < \sigma < \frac{\pi}{2},\ 0 \leqslant \beta < \sigma\}$ (these two families were originally studied by Karcher [13, 14]).



FIGURE 3. Left: $M_{\sigma,0,0}$, with $\sigma = \frac{\pi}{4}$. Right: $M_{\sigma,\alpha,0}$ for $\sigma = \alpha = \frac{\pi}{4}$.



FIGURE 4. $M_{\sigma,0,\beta}$, where $\sigma = \frac{\pi}{4}$ and $\beta = \frac{\pi}{8}$.

(1) When $\alpha = \beta = 0$, $M_{\sigma,0,0}$ contains four straight lines parallel to the x_1-axis, and $\mathrm{Iso}(M_{\sigma,0,0})$ is isomorphic to $(\mathbb{Z}/2\mathbb{Z})^4$ with generators S_1, S_2, S_3, R_D: S_1 is a reflection symmetry in a vertical plane orthogonal to the x_1-axis; S_2 is a reflection symmetry across a plane orthogonal to the x_2-axis; S_3 is a reflection symmetry in a horizontal plane (these three planes can be chosen meeting at a common point which is not contained in the surface); and R_D is the π-rotation around one of the four straight lines contained in the surface, see Figure 3 left. In this case, $T = (0, \pi\mu, 0)$.

(2) When $0 < \alpha < \frac{\pi}{2}$, $\mathrm{Iso}(M_{\sigma,\alpha,0})$ is isomorphic to $(\mathbb{Z}/2\mathbb{Z})^3$, with generators $\mathcal{D}$, S_2 and R_2, where S_2 represents a reflection symmetry across a plane orthogonal to the x_2-axis, and R_2 is a π-rotation around a line parallel to the x_2-axis that cuts $M_{\sigma,\alpha,0}$ orthogonally, see Figure 3 right. Now $T = (0, \pi\mu t_\alpha, 0)$, with $t_\alpha = \frac{\sin\sigma}{\sqrt{\sin^2\sigma\cos^2\alpha + \sin^2\alpha}}$.

(3) Suppose that $0 < \beta < \sigma$. Then $M_{\sigma,0,\beta}$ contains four straight lines parallel to the x_1-axis, and $\mathrm{Iso}(M_{\sigma,0,\beta})$ is isomorphic to $(\mathbb{Z}/2\mathbb{Z})^3$, with generators S_1, R_1 and R_D: S_1 represents a reflection symmetry across a plane orthogonal to the x_1-axis; R_1 corresponds to a π-rotation around a line parallel to the x_1-axis that cuts the surface orthogonally; and R_D is the π-rotation around any one of the straight lines contained in the surface, see Figure 4. Moreover, $T = (0, \pi\mu t^\beta, 0)$, where $t^\beta = \frac{\sin\sigma}{\sqrt{\sin^2\sigma - \sin^2\beta}}$.

Finally, it will be useful to see Σ_σ as a branched 2-covering of $\overline{\mathbb{C}}$ through the map $(z, w) \mapsto z$. Thus Σ_σ can be seen as two copies $\overline{\mathbb{C}}_1, \overline{\mathbb{C}}_2$ of $\overline{\mathbb{C}}$ glued along two common cuts γ_1, γ_2, which can be taken along the imaginary axis: γ_1 from D to D' and γ_2 from D'' to D'''.

5.1. **$M_{\sigma,\alpha,\beta}$ as a graph over $\{x_3 = 0\}/T$.** The KMR examples $M_{\sigma,\alpha,\beta}$ converge as $(\sigma, \alpha, \beta) \to (0, 0, 0)$ to a vertical catenoid, since Σ_σ converges to two pinched spheres, $g(z) \to z$ and $dh \to \pm\frac{dz}{z}$

as $(\sigma,\alpha,\beta)\to(0,0,0)$. In fact, we can obtain two catenoids in the limit, depending on the choice of branch for w (for each copy of $\overline{\mathbb{C}}$ in Σ_σ, we obtain one catenoid in the limit). One of our aims along this paper consists of gluing KMR examples $M_{\sigma,\alpha,0}$ or $M_{\sigma,0,\beta}$ near this catenoidal limit, to a convenient compact piece of a deformed CHM surface $M_k(\varepsilon/2)$. In this subsection we express part of $M_{\sigma,\alpha,\beta}$ as a vertical graph over the $\{x_3=0\}$ plane when σ,α,β are small.

Consider $M_{\sigma,\alpha,\beta}$ near the catenoidal limit, i.e. σ,α,β close to zero. Without lost of generality, we can assume $dh\sim\frac{dz}{z}$ in $\overline{\mathbb{C}}_1$. We are studying the surface in an annulus about one of its ends, say a zero of its Gauss map.

Lemma 5.2. *Consider $\alpha+\beta+\sigma\leqslant\varepsilon$ small. Up to translations, $M_{\sigma,\alpha,\beta}$ can be parametrized in the annulus $\{(z,w)\in\Sigma_\sigma \mid z\in\overline{\mathbb{C}}_1,\ |\frac{b}{a}|<|z|<\nu\}$ (for $\nu>|\frac{b}{a}|$ small) as:*

$$\begin{cases} X_1+i\,X_2=\frac{-1}{2}\left(z+\frac{1}{\overline{z}}\right)-\frac{(1+i)\overline{b}}{4\overline{z}^2}+\mathcal{O}(\varepsilon z^{-1}+\varepsilon^2 z^{-3}) \\ \\ X_3=\ln|z|+\mathcal{O}(\varepsilon^2 z^{-2}), \end{cases}$$

Proof. Recall we have assumed $dh\sim\frac{dz}{z}$ in the annulus we are working on. More precisely, we have

$$dh=\frac{\mu\,dz}{\sqrt{(z^2+\lambda^2)(z^2+\lambda^{-2})}}=\frac{\mu}{\lambda\sqrt{1+\lambda^{-2}z^2+\lambda^{-2}z^{-2}+\lambda^{-4}}}\,\frac{dz}{z}.$$

Since $\frac{\mu}{\lambda}=\frac{\pi}{(1+\cos(\sigma))\mathcal{K}(\sin^2\sigma)}=1+\mathcal{O}(\sigma^4)$, and $\lambda=\cot\frac{\sigma}{2}=\mathcal{O}(\varepsilon)$, we get

$$dh=\frac{dz}{z}(1+\mathcal{O}(\varepsilon^4))(1+\mathcal{O}(\varepsilon^2z^2+\varepsilon^2z^{-2}+\varepsilon^4)).$$

Since $|z|<\nu<1$, then

$$dh=\frac{dz}{z}(1+\mathcal{O}(\varepsilon^2z^{-2})).$$

Fix any point $z_0\in\mathbb{C}_1$, $z_0\notin\left\{\frac{-b}{a},\frac{\overline{a}}{\overline{b}}\right\}$ (which correspond to two ends of the KMR example), and recall that $g=\frac{az+b}{i(\overline{a}-\overline{b}z)}$. Straightforward computations give us, for $|b/a|<|z|<1$,

- $\int_{z_0}^{z}\frac{dh}{g}=-\frac{i\,\overline{b}}{a}\ln z-\frac{2i}{a^2z}+\frac{2i\,b}{a^3z^2}+C_1+\mathcal{O}(\varepsilon^2z^{-3})$,
- $\int_{z_0}^{z}g\,dh=-\frac{i\,b}{\overline{a}}\ln z-\frac{2i}{\overline{a}^2}z+C_2+\mathcal{O}(\varepsilon^2z^{-1})$,

where $C_1,C_2\in\mathbb{C}$ verify $\overline{C_1}-C_2=\frac{1}{2}\left(z_0+\frac{1}{\overline{z_0}}\right)+\mathcal{O}(\varepsilon)$. Taking into account that $a=(1+i)+\mathcal{O}(\varepsilon)$, we obtain

$$\begin{aligned} X_1+i\,X_2 &=\frac{1}{2}\left(\overline{\int_{z_0}^{z}\frac{dh}{g}}-\int_{z_0}^{z}g\,dh\right)\\ &=\frac{i}{\overline{a}^2}\left(z+\frac{1}{\overline{z}}\right)+\frac{i\,b}{\overline{a}}\ln|z|-\frac{i\,\overline{b}}{\overline{a}^3\overline{z}^2}+\frac{1}{2}\left(z_0+\frac{1}{\overline{z_0}}\right)+\mathcal{O}(\varepsilon^2z^{-3})\\ &=\frac{-1}{2}\left(z+\frac{1}{\overline{z}}\right)-\frac{(1+i)\overline{b}}{4\overline{z}^2}+\frac{1}{2}\left(z_0+\frac{1}{\overline{z_0}}\right)+\mathcal{O}(\varepsilon z^{-1}+\varepsilon^2z^{-3}). \end{aligned}$$

Similarly, $\int_{z_0}^{z}dh=\ln z-\ln z_0+\mathcal{O}(\varepsilon^2z^{-2})$, hence

$$X_3=\Re\int_{z_0}^{z}dh=\ln|z|-\ln|z_0|+\mathcal{O}(\varepsilon^2z^{-2}),$$

which completes the proof of lemma 5.2. □

By suitably translating $M_{\sigma,\alpha,\beta}$, we can assume its coordinate functions are as in Lemma 5.2.

Lemma 5.3. *Let (r,θ) denote the polar coordinates in the $\{x_3=0\}$ plane. If $\alpha+\beta+\sigma\leqslant\varepsilon$ small, then a piece of $M_{\sigma,\alpha,\beta}$ can be written as a vertical graph of the function*

$$\widetilde{U}(r,\theta)=-\ln(2r)+r\big(\kappa_1\cos\theta-\kappa_2\sin\theta\big)+\mathcal{O}(\varepsilon),$$

for $(r,\theta)\in(\frac{1}{4\sqrt{\varepsilon}},\frac{4}{\sqrt{\varepsilon}})\times[0,2\pi)$, where $\kappa_1=\Re(b)+\Im(b)$ and $\kappa_2=\Re(b)-\Im(b)$. We denote by $M_{\sigma,\alpha,\beta}(\gamma,\xi)$ the KMR example $M_{\sigma,\alpha,\beta}$ dilated by $1+\gamma$, for $\gamma\leqslant 0$ small, and translated by a vector $\xi=(\xi_1,\xi_2,\xi_3)$. Then, $M_{\sigma,\alpha,\beta}(\gamma,\xi)$ can be written as a vertical graph of

$$\widetilde{U}_{\gamma,\xi}(r,\theta)=-(1+\gamma)\ln\frac{2r}{1+\gamma}+r\left(\kappa_1\cos\theta-\kappa_2\sin\theta\right)-\frac{1+\gamma}{r}(\xi_1\cos\theta+\xi_2\sin\theta)+\xi_3+\mathcal{O}(\varepsilon).$$

Remark 5.4. *Recall that $b=\sin\frac{\alpha-\beta}{2}+i\sin\frac{\alpha+\beta}{2}$. In particular:*

- *When $\beta=0$, we have $\kappa_1=2\sin\frac{\alpha}{2}$ and $\kappa_2=0$, so*
$$\widetilde{U}_{\gamma,\xi}(r,\theta)=-(1+\gamma)\ln\frac{2r}{1+\gamma}+r\,\kappa_1\,\cos\theta-\frac{1+\gamma}{r}(\xi_1\cos\theta+\xi_2\sin\theta)+\xi_3+\mathcal{O}(\varepsilon).$$
- *When $\alpha=0$, $\kappa_1=0$ and $\kappa_2=2\sin\frac{\beta}{2}$, so*
$$\widetilde{U}_{\gamma,\xi}(r,\theta)=-(1+\gamma)\ln\frac{2r}{1+\gamma}-r\,\kappa_2\,\sin\theta-\frac{1+\gamma}{r}(\xi_1\cos\theta+\xi_2\sin\theta)+\xi_3+\mathcal{O}(\varepsilon).$$
- *When $\alpha=0$ (resp. $\beta=0$), we will consider $\xi_1=0$ (resp. $\xi_2=0$) in Section 7.*

Proof. Suppose $|\frac{b}{a}|<|z|<\nu$, $\nu>|\frac{b}{a}|$ small. From Lemma 5.2, we know the coordinate functions (X_1,X_2,X_3) of the perturbed KMR example $M_{\sigma,\alpha,\beta}(\gamma,\xi)$ are given by

$$\left\{\begin{array}{l} X_1+i\,X_2=-\frac{1+\gamma}{2}\left(z+\frac{1}{\overline{z}}\right)+A(z)\\[2ex] X_3=(1+\gamma)\ln|z|+\xi_3+\mathcal{O}(\varepsilon^2z^{-2}),\end{array}\right. \tag{23}$$

where

$$\begin{aligned}A(z)&=-\frac{(1+\gamma)(1+i)\overline{b}}{4\overline{z}^2}+(\xi_1+i\xi_2)+\mathcal{O}(\varepsilon z^{-1}+\varepsilon^2z^{-3})\\&=-\frac{(1+\gamma)(\kappa_1+i\kappa_2)}{4\overline{z}^2}+(\xi_1+i\xi_2)+\mathcal{O}(\varepsilon z^{-1}+\varepsilon^2z^{-3})\end{aligned}$$

If we set $z=|z|e^{i\psi}$ and $X_1+iX_2=re^{i\theta}$, then $z+\frac{1}{\overline{z}}=\left(|z|+\frac{1}{|z|}\right)e^{i\psi}$ and

$$\begin{aligned}r\cos\theta&=-\frac{1+\gamma}{2}\left(|z|+\frac{1}{|z|}\right)\cos\psi+A_1,\\ r\sin\theta&=-\frac{1+\gamma}{2}\left(|z|+\frac{1}{|z|}\right)\sin\psi+A_2,\end{aligned}$$

where $A_1=\Re(A)$ and $A_2=\Im(A)$. Therefore,

$$\begin{aligned}r^2=\;&\frac{(1+\gamma)^2}{4}\left(|z|+\frac{1}{|z|}\right)^2\left(1-\frac{4|z|}{(1+\gamma)(|z|^2+1)}\left(A_1\cos\psi+A_2\sin\psi\right)\right.\\&\left.+\frac{4|z|^2}{(1+\gamma)^2(|z|^2+1)^2}(A_1^2+A_2^2)\right).\end{aligned} \tag{24}$$

When $\frac{\sqrt{\varepsilon}}{4}\leqslant|z|\leqslant4\sqrt{\varepsilon}$, the functions A_i are bounded, and we get

$$r=\frac{1+\gamma}{2}\left(|z|+\frac{1}{|z|}\right)(1+\mathcal{O}(\sqrt{\varepsilon}))=\frac{1+\gamma}{2|z|}+\mathcal{O}(\sqrt{\varepsilon}). \tag{25}$$

In particular, $r = \mathcal{O}(1/\sqrt{\varepsilon})$. Moreover, we get $\frac{r}{\frac{1+\gamma}{2}\left(|z|+\frac{1}{|z|}\right)} = 1 + \mathcal{O}(\sqrt{\varepsilon})$, from where

$$\frac{X_1 + iX_2}{\frac{1+\gamma}{2}\left(|z| + \frac{1}{|z|}\right)} = e^{i\theta}(1 + \mathcal{O}(\sqrt{\varepsilon})).$$

On the other hand,

$$\frac{X_1 + iX_2}{\frac{1+\gamma}{2}\left(|z| + \frac{1}{|z|}\right)} = -e^{i\psi} + \frac{2|z|A}{(1+\gamma)(1+|z|^2)} = -e^{i\psi} + \mathcal{O}(\sqrt{\varepsilon}),$$

thus $e^{i\psi} = -e^{i\theta}(1 + \mathcal{O}(\sqrt{\varepsilon}))$.

From (24) and (25) we can deduce $\frac{(1+\gamma)^2(1+|z|^2)^2}{4|z|^2} = r^2\left(1 + \frac{2}{r}\left(A_1 \cos\psi + A_2 \sin\psi\right) + \mathcal{O}(\varepsilon)\right)$, from where we obtain

$$\begin{aligned} \frac{1}{|z|^2} &= \left(\frac{2r}{1+\gamma}\right)^2 \left(1 + \frac{2}{r}\left(A_1 \cos\psi + A_2 \sin\psi\right) + \mathcal{O}(\varepsilon)\right)(1 + \mathcal{O}(\varepsilon)) \\ &= \left(\frac{2r}{1+\gamma}\right)^2 \left(1 + \frac{2}{r}\left(A_1 \cos\psi + A_2 \sin\psi\right) + \mathcal{O}(\varepsilon)\right), \end{aligned} \tag{26}$$

and then

$$-\ln|z| = \ln\frac{2r}{1+\gamma} + \frac{1}{r}(A_1 \cos\psi + A_2 \sin\psi) + \mathcal{O}(\varepsilon). \tag{27}$$

Finally, it is not difficult to prove that

$$A_1 = -\frac{1+\gamma}{4|z|^2}\left(\kappa_1 \cos(2\psi) - \kappa_2 \sin(2\psi)\right) + \xi_1 + \mathcal{O}(\sqrt{\varepsilon})$$

$$\text{and} \qquad A_2 = -\frac{1+\gamma}{4|z|^2}\left(\kappa_1 \sin(2\psi) + \kappa_2 \cos(2\psi)\right) + \xi_2 + \mathcal{O}(\sqrt{\varepsilon}).$$

Therefore,

$$\begin{aligned} A_1 \cos\psi + A_2 \sin\psi &= -\frac{1+\gamma}{4|z|^2}\left(\kappa_1 \cos\psi - \kappa_2 \sin\psi\right) + \xi_1 \cos\psi + \xi_2 \sin\psi + \mathcal{O}(\sqrt{\varepsilon}) \\ &= -\frac{r^2}{1+\gamma}\left(\kappa_1 \cos\theta - \kappa_2 \sin\theta\right) + \xi_1 \cos\theta + \xi_2 \sin\theta + \mathcal{O}(\sqrt{\varepsilon}). \end{aligned}$$

From here, (27) and (23), Lemma 5.3 follows. □

5.2. **Parametrization of the KMR example on the cylinder.** In this subsection we want to parametrize the KMR example $M_{\sigma,\alpha,\beta}$ on a cylinder. Recall its conformal compactification Σ_σ only depends on σ. The parameter $\sigma \in (0, \frac{\pi}{2})$ will remains fixed along this subsection, and we will omit the dependence on σ of the functions we are introducing. Also recall that Σ_σ can be seen as a branched 2-covering of $\overline{\mathbb{C}}$, by gluing $\overline{\mathbb{C}}_1, \overline{\mathbb{C}}_2$ along two common cuts γ_1 and γ_2 along the imaginary axis joining the branch points D, D' and D'', D''' respectively (see (22)).

We introduce the sphero-conal coordinates (x, y) on the annulus $\mathbb{S}^2 - (\gamma_1 \cup \gamma_2)$ (see [11]): for any $(x, y) \in S^1 \times (0, \pi) \equiv [0, 2\pi) \times (0, \pi)$, we define

$$F(x, y) = (\cos x \, \sin y, \sin x \, m(y), l(x) \, \cos y) \in \mathbb{S}^2 - (\gamma_1 \cup \gamma_2),$$

where

$$m(y) = \sqrt{1 - \cos^2\sigma \cos^2 y} \quad \text{and} \quad l(x) = \sqrt{1 - \sin^2\sigma \sin^2 x}.$$

To understand the geometrical meaning of these variables, remark that $\{x = \mathit{constant}\}$ and $\{y = \mathit{constant}\}$ correspond to two closed curves on $\mathbb{S}^2$ which are two curves are the cross sections of the sphere and two elliptic cones (one with horizontal axis, the other one with vertical axis) having as vertex the center of the sphere.

If we compose $F(x,y)$ with the stereographical projection and enlarge the domain of definition of the function, we obtain the differentiable map $\mathbf{z}$ defined on the torus $S^1 \times S^1 \equiv [0,2\pi) \times [0,2\pi) \to \overline{\mathbb{C}}$ and given by

$$\mathbf{z}(x,y) = \frac{\cos x\,\sin y + i\,\sin x\, m(y)}{1 - l(x)\,\cos y}, \tag{28}$$

which is a branch 2-covering of $\overline{\mathbb{C}}$ with branch values in the four points whose sphero-conal coordinates are $(x,y) \in \left\{\pm\frac{\pi}{2}\right\} \times \{0,\pi\}$, which correspond to D, D', D'', D'''. Moreover, $\mathbf{z}$ maps $S^1 \times (0,\pi)$ on $\overline{\mathbb{C}} - (\gamma_1 \cup \gamma_2)$. Hence we can parametrize the KMR example by $\mathbf{z}$, via its Weierstrass data

$$g(\mathbf{z}) = \frac{a\mathbf{z}+b}{i(\overline{a} - \overline{b}\mathbf{z})}, \qquad dh = \mu\,\frac{d\mathbf{z}}{\sqrt{(\mathbf{z}^2 + {}^2)(\mathbf{z}^2 + {}^{-2})}},$$

We denote by $\widetilde{M}_{\sigma,\alpha,\beta}$ the lifting of $M_{\sigma,\alpha,\beta}$ to $\mathbb{R}^3/T$ by forgetting its non horizontal period (i.e. its period in homology, $\widetilde{T}$). We can then parametrize $\widetilde{M}_{\sigma,\alpha,\beta}$ on $S^1 \times \mathbb{R}$ by extending $\mathbf{z}$ to $[0,2\pi) \times \mathbb{R}$ periodically. But such a parametrization is not conformal, since the sphero-conal coordinates $(x,y) \mapsto F(x,y)$ of the sphere are not conformal. As the stereographic projection is a conformal map, it suffices to find new conformal coordinates (u,v) of the sphere defined on the cylinder. In particular, we look for a change of variables $(x,y) \mapsto (u,v)$ for which $|\widetilde{F}_u| = |\widetilde{F}_v|$ and $\langle \widetilde{F}_u, \widetilde{F}_v \rangle = 0$, where $\widetilde{F}(u,v) = F(x(u,v), y(u,v))$.

We observe that

$$|F_x| = \frac{\sqrt{k(x,y)}}{l(x)} \quad \text{and} \quad |F_y| = \frac{\sqrt{k(x,y)}}{m(y)},$$

with $k(x,y) = \sin^2\sigma\,\cos^2 x + \cos^2\sigma\,\sin^2 y$. Then it is natural to consider the change of variables $(x,y) \in [0,2\pi) \times \mathbb{R} \mapsto (u,v) \in [0,U_\sigma) \times \mathbb{R}$ defined by

$$u(x) = \int_0^x \frac{1}{l(t)}dt \quad \text{and} \quad v(y) = \int_{\frac{\pi}{2}}^y \frac{1}{m(t)}dt, \tag{29}$$

where

$$U_\sigma = u(2\pi) = \textstyle\int_0^{2\pi} \frac{dt}{\sqrt{1-\sin^2\sigma\,\sin^2 t}}. \tag{30}$$

Note that U_σ is a function on σ that goes to 2π as σ approaches to zero, and that the above change of variables is well defined because $\sigma \in (0,\frac{\pi}{2})$.

In these variables (u,v), $\mathbf{z}$ is v-periodic with period

$$V_\sigma = v(2\pi) - v(0) = \textstyle\int_0^{2\pi} \frac{dt}{\sqrt{1-\cos^2\sigma\,\cos^2 t}}. \tag{31}$$

The period V_σ goes to $+\infty$ as σ goes to zero (see lemma 5.5), which was clear taking into account the limits of $M_{\sigma,\alpha,\beta}$ as σ tends to zero.

From all this, we can deduce that $\widetilde{M}_{\sigma,\alpha,\beta}$ (resp. $M_{\sigma,\alpha,\beta}$) is conformally parametrized on $(u,v) \in I_\sigma \times \mathbb{R}$ (resp. $(u,v) \in I_\sigma \times J_\sigma$), where $I_\sigma = [0,U_\sigma]$ and $J_\sigma = [v(0), v(2\pi)]$.

In section 6, devoted to the study of the mapping properties of the Jacobi operator about $\widetilde{M}_{\sigma,\alpha,\beta}$, we will make exclusive use of the (u,v) variables. In the proof of Lemma 5.3 we have written as a vertical graph an appropriate piece of $\widetilde{M}_{\sigma,\alpha,\beta}$ corresponding to the annulus $\frac{\sqrt{\varepsilon}}{4} \leqslant |\mathbf{z}| \leqslant 4\sqrt{\varepsilon}$. The boundary curve along which we will glue the piece of $\widetilde{M}_{\sigma,\alpha,\beta}$ with the CHM surface corresponds to $|\mathbf{z}| = \sqrt{\varepsilon}$.

Lemma 5.5. *If* $\frac{\sqrt{\varepsilon}}{4} \leqslant |\mathbf{z}| \leqslant 4\sqrt{\varepsilon}$, *then*

$$-\frac{1}{2}\ln\varepsilon + c_1 \leqslant v \leqslant -\frac{1}{2}\ln\varepsilon + c_2,$$

where c_1 *and* c_2. *Under the same assumptions* $V_\sigma = -4\ln\varepsilon + \mathcal{O}(1)$.

Proof. The proof is articulated in two steps. Firstly, using equation (28), we will give the values of the y variable for which $\frac{\sqrt{\varepsilon}}{4} \leqslant |\mathbf{z}(x,y)| \leqslant 4\sqrt{\varepsilon}$ with $\sigma = \mathcal{O}(\varepsilon)$. Secondly we will determine the values of the v variable corresponding to these values of y. It is possible to show that, if $|\mathbf{z}|$ varies in the interval specified above, then $\pi - a_1 \leqslant y \leqslant \pi - a_2$, with $a_i = d_i\sqrt{\varepsilon}$, where $d_1 > d_2 > 0$ are constants. Being v an increasing function of y, then $v(\pi - a_1) \leqslant v(y) \leqslant v(\pi - a_2)$. Now we will obtain the values of $v(y)$ for $y = \pi - a_i$, $i = 1, 2$. To this aim we observe that

$$v(y) = \int_{\frac{\pi}{2}}^{y} \frac{ds}{\sqrt{1 - \cos^2\sigma\cos^2 s}}.$$

It holds that

$$\int_0^y \frac{ds}{\sqrt{1-\cos^2\sigma\cos^2 s}} = \int_0^y \frac{ds}{\sqrt{1-\cos^2\sigma + \cos^2\sigma\sin^2 s}} =$$

$$\frac{1}{\sin\sigma}\int_0^y \frac{ds}{\sqrt{1 + \frac{\cos^2\sigma}{\sin^2\sigma}\sin^2 s}} = \frac{1}{\sin\sigma}\mathcal{F}(y, m_\sigma),$$

where $m_\sigma = -\frac{\cos^2\sigma}{\sin^2\sigma}$ and $\mathcal{F}(y,m) = \int_0^y \frac{ds}{\sqrt{1-m\sin^2 s}}$ is the incomplete elliptic integral of the first kind. $\mathcal{F}(y,m)$ is an odd function with respect to y and if $k \in \mathbb{Z}$,

$$\mathcal{F}(y + k\pi, m) = \mathcal{F}(y,m) + 2k\mathcal{K}(m),$$

where $\mathcal{K}(m) = \mathcal{F}(\frac{\pi}{2}, m)$ is the complete elliptic integral of first kind. If $y = d\sqrt{\varepsilon}$ and $\sigma = \mathcal{O}(\varepsilon)$, then

$$\frac{1}{\sin\sigma}\mathcal{F}(y, m_\sigma) = -\frac{1}{2}\ln\varepsilon + \mathcal{O}(1).$$

It is known that if m is sufficiently big then

$$\mathcal{K}(m) = \frac{1}{\sqrt{-m}}\left(\ln 4 + \frac{1}{2}\ln(-m)\right)\left(1 + \mathcal{O}\left(\frac{1}{m}\right)\right).$$

It follows that

$$\frac{1}{\sin\sigma}\mathcal{K}(m_\sigma) = -\ln\varepsilon + \mathcal{O}(1).$$

Then, for $i = 1, 2$,

$$v(\pi - a_i) = \frac{1}{\sin\sigma}\left(\mathcal{F}(\pi - a_i, m_\sigma) - \mathcal{K}(m_\sigma)\right) =$$

$$\frac{1}{\sin\sigma}\left(\mathcal{F}(-a_i, m_\sigma) + 2\mathcal{K}(m_\sigma) - \mathcal{K}(m_\sigma)\right) =$$

$$\frac{1}{\sin\sigma}\left(-\mathcal{F}(d_i\sqrt{\varepsilon}, m_\sigma) + \mathcal{K}(m_\sigma)\right) = -\frac{1}{2}\ln\varepsilon + c_i.$$

The result concerning $V_\sigma = v(2\pi) - v(0)$ follows at once after having observed that $v(2\pi) = \frac{3\mathcal{K}(m_\sigma)}{\sin\sigma}$ and $v(0) = -\frac{\mathcal{K}(m_\sigma)}{\sin\sigma}$. □

From the lemma just proved it follows that the value of the v corresponding to $|\mathbf{z}| = \sqrt{\varepsilon}$, is $v_\varepsilon = -\frac{1}{2}\ln\varepsilon + c$, where c is a constant.

6. The Jacobi operator about KMR examples

The Jacobi operator for $M_{\sigma,\alpha,\beta}$ is given by $\mathcal{J} = \Delta_{ds^2} + |A|^2$, where $|A|^2$ is the squared norm of the second fundamental form on $M_{\sigma,\alpha,\beta}$ and Δ_{ds^2} is the Laplace-Beltrami operator with respect to the metric ds^2 induced on the surface by the immersion $X = (X_1, X_2, X_3)$ defined in section 5.1; i.e.

$$ds^2 = \frac{1}{4}\left(|g| + |g|^{-1}\right)^2 |dh|^2.$$

We consider the metric on the torus Σ_σ obtained as pull-back of standard metric ds_0^2 of the sphere $\mathbb{S}^2$ by the Gauss map N: $dN^*(ds_0^2) = -K\ ds^2$, where $K = -\frac{1}{2}|A|^2$ denotes the Gauss curvature of $M_{\sigma,\alpha,\beta}$. Hence $\Delta_{ds^2} = -K\,\Delta_{ds_0^2}$, and so

$$\mathcal{J} = -K\,(\Delta_{ds_0^2} + 2).$$

From [11] and taking into account the parametrization of $M_{\sigma,\alpha,\beta}$ on the cylinder given in subsection 5.2, we can deduce that, in the (x, y)-variables,

$$\Delta_{ds_0^2} := \frac{l(x)\,m(y)}{k(x,y)}\left[\partial_x\left(\frac{l(x)}{m(y)}\,\partial_x\right) + \partial_y\left(\frac{m(y)}{l(x)}\,\partial_y\right)\right].$$

In the (u, v)-variables defined by (29), we have $\mathcal{J} = \frac{-K}{k(x(u),y(v))}\mathcal{L}_\sigma$, where

$$\mathcal{L}_\sigma := \partial^2_{uu} + \partial^2_{vv} + 2\sin^2\sigma\,\cos^2(x(u)) + 2\cos^2\sigma\,\sin^2(y(v)) \tag{32}$$

is the Lamé operator [11].

Remark 6.1. *In Proposition 6.5, we will take $\sigma \to 0$. For such a limit, the torus Σ_σ degenerates into a Riemann surface with nodes consisting of two spheres jointed by two common points p_0, p_1, and the corresponding Jacobi operator equals the Legendre operator on $\mathbb{S}^2 \setminus \{p_0, p_1\}$, given by $\mathcal{L}_0 = \partial^2_{xx} + \sin y\,\partial_y\,(\sin y\,\partial_y) + 2\sin^2 y$ in the (x, y)-variables. Note that when $\sigma = 0$, the change of variables $(x, y) \mapsto (u, v)$ given in (29) is not well-defined.*

6.1. **The mapping properties of the Jacobi operator.**

From now on, we consider the conformal parametrization of $\widetilde{M}_{\sigma,\alpha,\beta}$ on the cylinder $\mathbb{S}^1 \times \mathbb{R} \equiv I_\sigma \times \mathbb{R}$ described in subsection 5.2. Our aim along this subsection is to study the mapping properties of the operator $\mathcal{J}$. It is clear that it is sufficient to study the simpler operator $\mathcal{L}_\sigma$ defined by (32), so we will study the problem

$$\begin{cases} \mathcal{L}_\sigma w = f & \text{in } I_\sigma \times [v_0, +\infty[\\ w = \varphi & \text{on } I_\sigma \times \{v_0\} \end{cases}$$

with $v_0 \in \mathbb{R}$ and considering convenient normed functional spaces for w, f, φ so that the norm of w is bounded by the one of f.

We will work in two different functional spaces to solve the above Dirichlet problem. To explain the reason, it is convenient to recall that the isometry group of $\widetilde{M}_{\sigma,\alpha,\beta}$ depends on the values of the three parameters σ, α, β. When $\alpha = 0$ (resp. $\beta = 0$), $\widetilde{M}_{\sigma,\alpha,\beta}$ is invariant by the reflection symmetry about the $\{x_1 = 0\}$ plane (resp. $\{x_2 = 0\}$ plane). We are interested in showing the existence of families of minimal surfaces close to $\widetilde{M}_{\sigma,0,\beta}$ and $\widetilde{M}_{\sigma,\alpha,0}$, keeping the same properties of symmetry. Thus the surfaces in the family about $\widetilde{M}_{\sigma,0,\beta}$ (resp. $\widetilde{M}_{\sigma,\alpha,0}$) will be defined as normal graphs of functions defined in $I_\sigma \times \mathbb{R}$ which are even (resp. odd) in the first variable. We will solve the above Dirichlet problem in the first case. The second one follows similarly.

Definition 6.2. *Given $\sigma \in (0,\pi/2)$, $\ell \in \mathbb{N}$, $\alpha \in (0,1)$, $\mu \in \mathbb{R}$ and an interval I, we define $\mathcal{C}^{\ell,\alpha}_{\mu}(I_\sigma \times I)$ as the space of functions $w = w(u,v)$ in $\mathcal{C}^{\ell,\alpha}_{loc}(I_\sigma \times I)$ which are even and U_σ-periodic in the variable u and for which the following norm is finite:*

$$\|w\|_{\mathcal{C}^{\ell,\alpha}_{\mu}(I_\sigma\times I)} := \sup_{v\in I} e^{-\mu v}\|w\|_{\mathcal{C}^{\ell,\alpha}(I_\sigma\times[v,v+1])}$$

It is possible to prove that $U_\sigma \to 2\pi$ as $\sigma \to 0$. We observe that the Jacobi operator $\mathcal{L}_\sigma$ becomes a Fredholm operator when restricted to the functional spaces defined in 6.2. Moreover, $\mathcal{L}_\sigma$ has separated variables. Then we consider the operator

$$L_\sigma = \partial^2_{uu} + 2\sin^2\sigma\cos^2(x(u)),$$

defined on the space of U_σ-periodic and even functions in I_σ. This operator L_σ is uniformly elliptic and self-adjoint. In particular, L_σ has discrete spectrum $(\lambda_{\sigma,i})_{i\geqslant 0}$, that we assume arranged so that $\lambda_{\sigma,i} < \lambda_{\sigma,i+1}$ for every i. Each eigenvalue $\lambda_{\sigma,i}$ is simple because we only consider even functions. We denote by $e_{\sigma,i}$ the even eigenfunction associated to $\lambda_{\sigma,i}$, normalized so that

$$\int_0^{U_\sigma} (e_{\sigma,i}(u))^2\, du = 1.$$

Lemma 6.3. *For every $i \geqslant 0$, the eigenvalue $\lambda_{\sigma,i}$ of the operator L_σ and its associated eigenfunctions $e_{\sigma,i}$ satisfy*

$$-2\sin^2\sigma \leqslant \lambda_{\sigma,i} - i^2 \leqslant 0, \qquad |e_{\sigma,i} - e_{0,i}|_{\mathcal{C}^2} \leqslant c_i \sin^2\sigma, \tag{33}$$

where $e_{0,i}(u) := \cos(ix(u))$ for every $u \in I_\sigma$, and the constant $c_i > 0$ depends only on i (it does not depend on σ).

Proof. The bound for $\lambda_{\sigma,i} - i^2$ comes from the variational characterization of the eigenvalues

$$\lambda_{\sigma,i} = \sup_{\operatorname{codim} E = i} \inf_{\substack{e\in E \\ ||e||_{L^2}=1}} \int_0^{U_\sigma} \left((\partial_u e)^2 - 2\sin^2\sigma\cos^2(x(u))\, e^2\right) du,$$

where E is a subset of the space of U_σ-periodic and even functions in $L^2(I_\sigma)$, since it always holds $0 \leqslant 2\sin^2\sigma\cos^2(x(u)) \leqslant 2\sin^2\sigma$. The bound for the eigenfunctions follows from standard perturbation theory [15]. □

The Hilbert basis $\{e_{\sigma,i}\}_{i\in\mathbb{N}}$ of the space of U_σ-periodic and even functions in $L^2(I_\sigma)$ introduced above induces the following Fourier decomposition of functions $g = g(u,v)$ in $L^2(I_\sigma \times \mathbb{R})$ which are U_σ-periodic and even in the variable u,

$$g(u,v) = \sum_{i\geqslant 0} g_i(v)\, e_{\sigma,i}(u).$$

From this, we deduce that the operator $\mathcal{L}_\sigma$, can be decomposed as $\mathcal{L}_\sigma = \sum_{i\geqslant 0} L_{\sigma,i}$, being

$$L_{\sigma,i} = \partial^2_{vv} + 2\cos^2\sigma\sin^2(y(v)) - \lambda_{\sigma,i}\,, \quad \text{for every } i \geqslant 0.$$

Since $0 \leqslant 2\cos^2\sigma\sin^2(y(v)) \leqslant 2\cos^2\sigma = 2 - 2\sin^2\sigma$, Lemma 6.3 gives us

$$P_{\sigma,i} := 2\cos^2\sigma\,\sin^2(y(v)) - \lambda_{\sigma,i} \leqslant 2 - i^2. \tag{34}$$

This fact allows us to prove the following lemma, which assures that $\mathcal{L}_\sigma$ is injective when restricted to the set of functions that are, in the variable u, even and L^2-orthogonal to $e_{\sigma,0}$ and $e_{\sigma,1}$.

Lemma 6.4. *Given $v_0 < v_1$, let w be a solution of $\mathcal{L}_\sigma w = 0$ on $I_\sigma \times [v_0, v_1]$ such that*

(i) $w(\cdot\,, v_0) = w(\cdot\,, v_1) = 0$

(ii) $\int_0^{U_\sigma} w(u,v)e_{\sigma,i}(u)\,du = 0$ *, for every* $v \in [v_0, v_1]$ *and* $i \in \{0,1\}$

Then $w = 0$.

Proof. By *(ii)*, $w = \sum_{i \geqslant 2} w_i(v)\, e_{\sigma,i}(u)$. Since the potential $P_{\sigma,i}$ of the operator $L_{\sigma,i}$ is negative for every $i \geqslant 2$ (see (34)) and the operator $L_{\sigma,i}$ is elliptic, the maximum principle holds. We can then conclude the Lemma 6.4 from *(i)*. □

To study the mapping properties of the Jacobi operator $\mathcal{L}_\sigma$, we need to give a description of the Jacobi fields associated to $M_{\sigma,\alpha,0}$, which are defined as the solutions of $\mathcal{L}_\sigma v = 0$.

Since $M_{\sigma,\alpha,0}$ is invariant by the reflection symmetry across the plane $\{x_2 = 0\}$, there are only four independent Jacobi fields:

- Two Jacobi fields induced by vertical translations and by horizontal translations in the x_1-direction. These Jacobi fields are clearly periodic and hence bounded.
- A third Jacobi field generated by the 1-parameter group of dilations, which is not bounded (it grows linearly).
- The last Jacobi field can be obtained by considering the 1-parameter family of minimal surfaces induced by changing the parameter σ. This Jacobi field is not periodic and grows linearly.

The Jacobi fields induced by translation along the x_3-axis and by dilatation, are solutions of $\mathcal{L}_\sigma u = 0$ that are collinear to the eigenfunction $e_{\sigma,0}$. The Jacobi fields induced by the horizontal translation and by the variation of the parameter σ, are collinear to $e_{\sigma,1}$.

The Jacobi fields of $M_{\sigma,0,\beta}$, which is invariant by the reflection symmetry across the plane $\{x_1 = 0\}$ are the same as $M_{\sigma,\alpha,0}$, with the exception of that one induced by horizontal translations in the x_1-direction which is to be replaced by the field induced by horizontal translations in the x_2-direction.

The following proposition states that, if the parameter μ is appropriately chosen, there exists a right inverse for $\mathcal{L}_\sigma$ whose norm is uniformly bounded.

Proposition 6.5. *Given* $\mu \in (-2,-1)$*, there exists* $\sigma_0 \in (0, \pi/2)$ *so that, for every* $\sigma \in (0, \sigma_0)$ *and* $v_0 \in \mathbb{R}$*, there exists an operator*

$$G_{\sigma,v_0} : \; \mathcal{C}^{0,\alpha}_\mu(I_\sigma \times [v_0, +\infty)) \quad \longrightarrow \quad \mathcal{C}^{2,\alpha}_\mu(I_\sigma \times [v_0, +\infty))$$

such that for all $f \in \mathcal{C}^{0,\alpha}_\mu(I_\sigma \times [v_0, +\infty))$*, the function* $w := G_{\sigma,v_0}(f)$ *solves*

$$\begin{cases} \mathcal{L}_\sigma\, w = f & , \text{ on } I_\sigma \times [v_0, +\infty) \\ w \in \operatorname{Span}\{e_{\sigma,0}, e_{\sigma,1}\}, & \text{on } I_\sigma \times \{v_0\}. \end{cases} \tag{35}$$

Moreover $||w||_{\mathcal{C}^{2,\alpha}_\mu} \leqslant c\,||f||_{\mathcal{C}^{0,\alpha}_\mu}$*, for some constant* $c > 0$ *which does not depend on* $\sigma \in (0, \sigma_0)$ *nor* $v_0 \in \mathbb{R}$.

Proof. Every $f \in \mathcal{C}^{0,\alpha}_\mu(I_\sigma \times [v_0, +\infty))$ can be decomposed as

$$f = f_0\, e_{\sigma,0} + f_1\, e_{\sigma,1} + \bar{f},$$

where $\bar{f}(\cdot, v)$ is L^2-orthogonal to $e_{\sigma,0}$ and to $e_{\sigma,1}$, for each $v \in [v_0, +\infty)$.

Step 1. Firstly, let's prove Proposition 6.5 for functions $\bar{f} \in \mathcal{C}^{0,\alpha}_\mu(I_\sigma \times [v_0, +\infty))$ that are L^2-orthogonal to $\{e_{\sigma,0}, e_{\sigma,1}\}$. As a consequence of the Lemma 6.4, $\mathcal{L}_\sigma$ is injective when acting on the space of these functions. Hence, Fredholm alternative assures there exists, for each $v_1 > v_0 + 1$, a unique $\bar{w} \in \mathcal{C}^{2,\alpha}_\mu$, with $\bar{w}(\cdot, v)$ L^2-orthogonal to $e_{\sigma,0}, e_{\sigma,1}$ satisfying:

$$\begin{cases} \mathcal{L}_\sigma\, \bar{w} = \bar{f} & \text{on } I_\sigma \times [v_0, v_1], \\ \bar{w}(\cdot, v_0) = \bar{w}(\cdot, v_1) = 0. \end{cases} \tag{36}$$

Assertion 6.6. *There exist $c \in \mathbb{R}$ and $\sigma_0 \in (0, \pi/2)$ such that, for every $\sigma \in (0, \sigma_0)$, $v_0 \in \mathbb{R}$, $v_1 > v_0 + 1$ and $\bar{f} \in \mathcal{C}^{0,\alpha}_\mu(I_\sigma \times [v_0, v_1])$, there exists $\bar{w} \in \mathcal{C}^{2,\alpha}_\mu(I_\sigma \times [v_0, v_1])$ L^2-orthogonal to $\{e_{\sigma,0}, e_{\sigma,1}\}$ satisfying (36) and*

$$\sup_{I_\sigma \times [v_0,v_1]} e^{-\mu v} |\bar{w}| \leqslant c \sup_{I_\sigma \times [v_0,v_1]} e^{-\mu v} |\bar{f}|. \tag{37}$$

Suppose by contradiction that Assertion 6.6 is false. Then, for every $n \in \mathbb{N}$, there exists $\sigma_n \in (0, 1/n)$, $v_{1,n} > v_{0,n} + 1$ and $\bar{f}_n, \bar{w}_n$ satisfying (36) (for $\sigma_n, v_{0,n}, v_{1,n}$ instead of σ, v_0, v_1) such that

$$\sup_{I_{\sigma_n} \times [v_{0,n}, v_{1,n}]} e^{-\mu v} |\bar{f}_n| = 1, \qquad \text{and}$$

$$A_n := \sup_{I_{\sigma_n} \times [v_{0,n}, v_{1,n}]} e^{-\mu v} |\bar{w}_n| \to +\infty \quad \text{as} \quad n \to \infty.$$

Since $I_{\sigma_n} \times [v_{0,n}, v_{1,n}]$ is a compact set, A_n is achieved at a point $(u_n, v_n) \in I_{\sigma_n} \times [v_{0,n}, v_{1,n}]$.

After passing to a subsequence, the intervals $I_n = [v_{0,n} - v_n, v_{1,n} - v_n]$ converge to a set I_∞. Elliptic estimates imply that

$$\sup_{I_{\sigma_n} \times [v_{0,n}, v_{0,n}+\frac{1}{2}]} e^{-\mu v} |\nabla \bar{w}_n| \leqslant c\Big(\sup_{I_{\sigma_n} \times [v_{0,n}, v_{0,n}+1]} e^{-\mu v} |\bar{f}_n| + \sup_{I_{\sigma_n} \times [v_{0,n}, v_{0,n}+1]} e^{-\mu v} |\bar{w}_n|\Big) \leqslant c(1 + A_n).$$

From this estimate for the gradient of $\bar{w}_n$ near $v = v_{0,n}$ it follows that v_n cannot be too close to $v_{0,n}$, where $\bar{w}_n$ vanishes. More precisely, $v_{0,n} - v_n$ remains bounded away from 0, and then converges to some $\bar{v}_0 \in [-\infty, 0)$. Using similar arguments, it is possible to show that $\nabla \bar{w}_n$ is bounded near $v_{1,n}$ and consequently $v_{1,n} - v_n$ converges to some $\bar{v}_1 \in (0, \infty]$. Then we can conclude that $I_\infty = [\bar{v}_0, \bar{v}_1]$.

We define

$$\tilde{w}_n(u, v) := \frac{e^{-\mu v_n}}{A_n} \bar{w}_n(u, v + v_n), \quad (u, v) \in I_{\sigma_n} \times I_n.$$

We observe that

$$|\tilde{w}_n(u, v)| \leqslant e^{\mu\, v} \, \frac{e^{-\mu(v+v_n)} |\bar{w}_n(u, v + v_n)|}{A_n} \leqslant e^{\mu\, v}$$

$$\text{and} \quad \sup_{I_{\sigma_n} \times I_n} e^{-\mu v} |\tilde{w}_n| = 1. \tag{38}$$

Using the above estimate for $\nabla \bar{w}_n$ we obtain

$$|\nabla \tilde{w}_n| \leqslant c \frac{1 + A_n}{A_n} \, e^{\mu\, v} < 2c\, e^{\mu\, v}.$$

Since the sequences $\{\tilde{w}_n\}_n$ and $\{\nabla \tilde{w}_n\}_n$ are uniformly bounded, Ascoli-Arzelà's theorem assures that, if $n \to +\infty$ a subsequence of $\{\tilde{w}_n\}_n$ converges on compact sets of $I_0 \times I_\infty$ to a function $\tilde{w}_\infty$ which vanishes on $I_0 \times \partial I_\infty$, when $\partial I_\infty \neq \emptyset$, and such that $\tilde{w}_\infty(\cdot, v)$ is L^2-orthogonal to $\{e_{0,0}, e_{0,1}\}$ for each $v \in I_\infty$. Moreover,

$$\sup_{I_0 \times I_\infty} e^{-\mu v} |\tilde{w}_\infty| = 1. \tag{39}$$

Since $\sigma_n \to 0$ as $n \to \infty$, we can conclude that $\tilde{w}_\infty$ satisfies

$$\begin{cases} \mathcal{L}_0 \tilde{w}_\infty = 0 \\ \tilde{w}_\infty|_{I_0 \times \partial I_\infty} = 0 \text{ (if } \partial I_\infty \neq \emptyset) \end{cases}$$

If I_∞ is bounded, the maximum principle allows us to conclude that $\tilde{w}_\infty = 0$ on $I_0 \times I_\infty$, which contradicts (39). Hence I_∞ is an unbounded interval.

Recall $\mathcal{L}_0$ is given in terms of the (x,y)-variables. The equation $\mathcal{L}_0\tilde{w}_\infty = 0$ becomes

$$\partial^2_{xx}\tilde{w}_\infty + \sin y\,\partial_y\left(\sin y\,\partial_y\tilde{w}_\infty\right) + 2\sin^2 y\,\tilde{w}_\infty = 0.$$

Now we consider the eigenfunctions decomposition of $\tilde{w}_\infty$,

$$\tilde{w}_\infty(x,y) = \sum_{j\geqslant 2} a_j(y)\,\cos(j\,x).$$

Each coefficient a_j, $j\geqslant 2$, must satisfy the associated Legendre differential equation (see Appendix C)

$$\sin y\,\partial_y\left(\sin y\,\partial_y a_j\right) - j^2 a_j + 2\sin^2 y\,a_j = 0.$$

We obtain that $a_j(y)$ equals the associated Legendre functions of second kind, i.e. $a_j(y) = Q^j_1(\cos y)$, $j\geqslant 2$.

We obtain from (29) that

$$u(x)\to x \quad\text{and}\quad v(y)\to \frac{1}{2}\ln\left|\tan\frac{y}{2}\right| \quad \text{as } \sigma\to 0.$$

In particular, define $y(v) = 2\arctan(e^{2v})$ for $\sigma = 0$. Therefore,

$$\cos y(v) = \frac{1-e^{4v}}{1+e^{4v}} \qquad \text{and} \tag{40}$$

$$\tilde{w}_\infty(u,v) = \sum_{j\geqslant 2} Q^j_1\left(\frac{1-e^{4v}}{1+e^{4v}}\right)\cos(j\,u).$$

It is possible to show that $|a_j|$ tend to $+\infty$ as the function $e^{2j|v|}$. Since the interval I_∞ is unbounded, we reach a contradiction with (39). This proves Assertion 6.6.

Let $c\in\mathbb{R}$ and σ_0 satisfy Assertion 6.6 and fix $\sigma\in(0,\sigma_0)$, $v_0\in\mathbb{R}$ and $\bar{f}\in\mathcal{C}^{0,\alpha}_\mu(I_\sigma\times[v_0,v_1])$. Then, for every $v_1 > v_0+1$, there exists a function $\bar{w}$ which is L^2-orthogonal to $\{e_{\sigma,0}, e_{\sigma,1}\}$ and satisfies (36) and (37). Let's take the limit as $v_1\to\infty$. Clearly,

$$e^{-\mu\,v}|\bar{w}| \leqslant \|\bar{w}\|_{\mathcal{C}^{0,\alpha}_\mu(I_\sigma\times[v_0,v_1])} \leqslant c\,\|\bar{f}\|_{\mathcal{C}^{0,\alpha}_\mu(I_\sigma\times[v_0,v_1])}.$$

And using Schäuder estimates we get

$$e^{-\mu\,v}|\nabla\bar{w}| \leqslant \|\bar{w}\|_{\mathcal{C}^{2,\alpha}_\mu(I_\sigma\times[v_0,v_1])} \leqslant c_1\left(\|\bar{f}\|_{\mathcal{C}^{0,\alpha}_\mu(I_\sigma\times[v_0,v_1])} + \|\bar{w}\|_{\mathcal{C}^{0}_\mu(I_\sigma\times[v_0,v_1])}\right) \leqslant c_2\|\bar{f}\|_{\mathcal{C}^{0,\alpha}_\mu(I_\sigma\times[v_0,v_1])}.$$

Hence Ascoli-Arzelà's theorem assures that a subsequence of $\{\bar{w}_{v_1}\}_{v_1>v_0+1}$ converges to a function $\bar{w}$ defined on $I_\sigma\times[v_0,+\infty)$, which satisfies

$$\sup_{I_\sigma\times[v_0,+\infty)} e^{-\mu\,v}|\bar{w}| \leqslant c \sup_{I_\sigma\times[v_0,+\infty)} e^{-\mu\,v}|\bar{f}|.$$

Using again elliptic estimates we can conclude that $\bar{w}$ satisfies the statement of Proposition 6.5. The uniqueness of the solution follows from Lemma 6.4.

Step 2 Let's now consider $f\in\mathcal{C}^{0,\alpha}_\mu(I_\sigma\times[v_0,+\infty))$ in $\mathrm{Span}\{e_{\sigma,0}, e_{\sigma,1}\}$, i.e.

$$f(u,v) = f_0(v)\,e_{\sigma,0}(u) + f_1(v)\,e_{\sigma,1}(u).$$

We extend the functions $f_0(v), f_1(v)$ for $v\leqslant v_0$ by $f_0(v_0), f_1(v_0)$, respectively. Given $v_1 > v_0+1$, consider the problem

$$\begin{cases} L_{\sigma,j}w_j = f_j, & v\in(-\infty,v_1] \\ w_j(v_1) = \partial_v w_j(v_1) = 0. \end{cases} \tag{41}$$

Peano theorem assures the existence and the uniqueness of the solution w_j. Our aim consists of proving the following result.

Assertion 6.7. *$\sup_{(-\infty,v_1]} e^{-\mu v}|w_j| \leqslant c \sup_{(-\infty,v_1]} e^{-\mu v}|f_j|$ for some constant c which does not depend on v_1.*

Suppose by contradiction that, for every $n \in \mathbb{N}$, there exists $\sigma_n \in (0, 1/n)$, $v_{1,n} > v_{0,n} + 1$ and $f_{j,n}, w_{j,n}$ satisfying (41) such that

$$\sup_{(-\infty,v_{1,n}]} e^{-\mu v}\,|\bar{f}_{j,n}| = 1,$$

$$A_n := \sup_{(-\infty,v_{1,n}]} e^{-\mu v}\,|w_{j,n}| \to +\infty \quad \text{as} \quad n \to \infty.$$

The solution $w_{j,n}$ of the previous equation is a linear combination of the two solutions of the homogeneous problem $L_{\sigma_n,j}\, w = 0$. They grow at most linearly at ∞ (recall that the Jacobi fields have this rate of growth). Hence the supremum A_n is achieved at a point $v_n \in (-\infty, v_{1,n}]$. We define on $I_n := (-\infty, v_{1,n} - v_n]$ the function

$$\tilde{w}_{j,n}(v) := \frac{1}{A_n}\, e^{-\mu v_n}\, w_{j,n}(v_n + v).$$

As in Step 1, one shows that the sequence $\{v_{1,n} - v_n\}_n$ remains bounded away from 0 and, after passing to a subsequence, it converges to $\bar{v}_1 \in (0, +\infty]$ and $\{\tilde{w}_{j,n}\}_n$ converges on compact subsets of $I_\infty = (-\infty, \bar{v}_1]$ to a nontrivial function $\tilde{w}_j$ such that

$$\sup_{v \in I_\infty} e^{-\mu v}\,|\tilde{w}_j| = 1 \tag{42}$$

and $\tilde{w}_j(\bar{v}_1) = \partial_v w_j(\bar{v}_1) = 0$, if $\bar{v}_1 < +\infty$. The function $\tilde{w}_j$ is the solution of a second order ordinary differential equation, given, in terms of the (x, y)-variables, by

$$\sin y\, \partial_y\,(\sin y\, \partial_y \tilde{w}_j) - j^2\, \tilde{w}_j + 2\sin^2 y\, \tilde{w}_j = 0. \tag{43}$$

If $\bar{v}_1 < +\infty$ then $\tilde{w}_j = 0$ and this is a contradiction with (42). In the case $\bar{v}_1 = +\infty$ we will try to reach a contradiction determining the solution (43). This is again the associated Legendre differential equation (see Appendix C). The solutions of the equation (43) are a linear combination of the associated Legendre functions of first kind, $P_1^j(\cos y)$, and second kind, $Q_1^j(\cos y)$, with $j = 0, 1$. It holds that $P_1^0(\cos y) = \cos y$ and $P_1^1(\cos y) = -\sin y$. We change of coordinates to express $\tilde{w}_j$ in terms of the (u, v)-variables. It is possible to show that as $v \to \pm\infty$ then $|Q_1^1(\cos y(v))|$ and $|Q_1^0(\cos y(v))|$ tend to ∞ respectively as $e^{2|v|}$ and $|v|$. We can conclude that functions $\tilde{w}_1$ and $\tilde{w}_0$ do not satisfy the equation (42) with $\mu \in (-2, -1)$, a contradiction.

Therefore, $\sup_{(-\infty,v_1]} e^{-\mu v}\,|w_j| \leqslant c\, \sup_{(-\infty,v_1]} e^{-\mu v}\,|f_j|$. Taking $v_1 \to +\infty$, we get a solution of $L_{\sigma,j}\, w_j = f_j$ defined in $[v_0, +\infty)$ which satisfies

$$\sup_{[v_0,+\infty)} e^{-\mu v}\,|w_j| \leqslant c \sup_{[v_0,+\infty)} e^{-\mu v}\,|f_j|.$$

Elliptic estimates allow us to obtain the wanted estimates for the derivatives. To prove the uniqueness of solution, it is sufficient to observe that no solution of $\mathcal{L}_\sigma v = 0$ that is collinear with $e_{\sigma,0}$ and $e_{\sigma,1}$ decays exponentially at ∞. This fact follows from the behaviour of the Jacobi fields. □

7. A FAMILY OF MINIMAL SURFACES CLOSE TO $\widetilde{M}_{\sigma,0,\beta}$ AND $\widetilde{M}_{\sigma,\alpha,0}$

The aim of this section is to find a family of minimal surfaces near to a translated and dilated copy of $\widetilde{M}_{\sigma,0,\beta}$ and $\widetilde{M}_{\sigma,\alpha,0}$ with given Dirichlet data on the boundary. We start recalling that in subsection 5.1 we observed that a translated and dilated copies of $\widetilde{M}_{\sigma,\alpha,0}$ and $\widetilde{M}_{\sigma,0,\beta}$ can be expressed as the graphs over the $\{x_3 = 0\}$ plane respectively of the functions

$$\tilde{U}_{\gamma,\xi}(r,\theta) = -(1+\gamma)\ln(2r) + r\kappa_1\cos\theta - \frac{1+\gamma}{r}\xi_1\cos\theta + d + \mathcal{O}(\varepsilon) \tag{44}$$

$$\tilde{U}_{\gamma,\xi}(r,\theta) = -(1+\gamma)\ln(2r) - r\kappa_2\sin\theta - \frac{1+\gamma}{r}\xi_2\sin\theta + d + \mathcal{O}(\varepsilon) \tag{45}$$

where $d = \xi_3 + (1+\gamma)\ln(1+\gamma)$, $\kappa_1, \kappa_2, \xi_1, \xi_2, \xi_3, \gamma \in \mathbb{R}$ small enough, $\kappa_1 = b_1 + b_2$, $\kappa_2 = b_1 - b_2$, $b_1 = \sin\frac{\alpha-\beta}{2}$, $b_2 = \sin\frac{\alpha+\beta}{2}$, and r belongs to a neighbourhood of $r_\varepsilon = \frac{1}{2\sqrt{\varepsilon}}$.

We denote by Z the immersion of the surface $\widetilde{M}_{\sigma,\alpha,\beta}$. The following proposition, whose proof is contained in section 10, states that the linearized of the mean curvature operator is the Lamé operator introduced in section 5.2.

Proposition 7.1. *The surface parameterized by $Z_f := Z + f\,N$ is minimal if and only if the function f is a solution of*

$$\mathcal{L}_\sigma f = Q_\sigma(f).$$

where $\mathcal{L}_\sigma := \partial^2_{uu} + \partial^2_{vv} + 2\sin^2\sigma\,\cos^2(x(u)) + 2\cos^2\sigma\,\sin^2(y(v))$ is the Lamé operator and Q_σ is a nonlinear operator which satisfies

$$\|Q_\sigma(f_2) - Q_\sigma(f_1)\|_{\mathcal{C}^{0,\alpha}(I_\sigma\times[v,v+1])} \leqslant c \sup_{i=1,2}\|f_i\|_{\mathcal{C}^{2,\alpha}(I_\sigma\times[v,v+1])}\,\|f_2 - f_1\|_{\mathcal{C}^{2,\alpha}(I_\sigma\times[v,v+1])} \tag{46}$$

for all f_1, f_2 such that $\|f_i\|_{\mathcal{C}^{2,\alpha}(I_\sigma\times[v,v+1])} \leqslant 1$. Here the constant $c > 0$ does not depend on $v \in \mathbb{R}$, nor on $\sigma \in (0, \frac{\pi}{2})$.

As a consequence of the dilation of factor $1+\gamma$ of the surface the minimal surface equation becomes

$$\mathcal{L}_\sigma\, w = \frac{1}{1+\gamma} Q_\sigma\left((1+\gamma)w\right). \tag{47}$$

We now truncate the surfaces $\widetilde{M}_{\sigma,\alpha,0}$ and $\widetilde{M}_{\sigma,0,\beta}$ at the graph of the curve $r = \frac{1}{2\sqrt{\varepsilon}}$ of the function (44) and (45) with, respectively, $\beta = 0$ and $\alpha = 0$ and we consider only the upper half of these surfaces which we call M_1 and M_2.

We are interested in minimal normal graphs over these surfaces which are asymptotic to them. The normal graph of the function w over M_1, M_2 is minimal, if and only if w is a solution of (47).

In subsection 5.1 we have proved lemma 5.3, which allows us to parametrize the surfaces M_1, M_2 in a neighbourhood of r_ε as the graph of the functions given above. The surfaces M_1 and M_2 are parametrized on $I_\sigma \times [v_\varepsilon, V_\sigma]$ where v_ε denote value of the variable v corresponding to the value r_ε of the variable r. In section 5 we showed that $v_\varepsilon = -\frac{1}{2}\ln\varepsilon + \mathcal{O}(1)$.

It is important to remark that the boundary of these surfaces does not correspond to the curve $v = v_\varepsilon$. We therefore modify the above parametrization so that for $v \in [v_\varepsilon - \ln 2, v_\varepsilon + \ln 2]$ the image of the functions (44) and (45) corresponds to the horizontal curve $v = v_\varepsilon$. Furthermore we would like that the normal vector field relative to M_1, M_2 is vertical near the boundary of this surface. This can be achieved by modifying the normal vector field into a transverse vector field $\tilde{N}$ which agrees with the normal vector field N for all $v \geqslant v_\varepsilon + \ln 4$ and with the vector e_3 for all $v \in [v_\varepsilon, v_\varepsilon + \ln 2]$.

Now, we consider a graph over this surface for some function u, using the modified vector field $\tilde{N}$. This graph will be minimal if and only if the function u is a solution of a nonlinear elliptic equation related to (47). To get the new equation, we take into account the effects of the change of parameterization and the change in the vector field N into $\tilde{N}$. The new minimal surface equation is

$$\mathcal{L}_\sigma\, w = \tilde{L}_\varepsilon\, w + \tilde{Q}_\sigma\,(w)\,. \tag{48}$$

Here $\tilde{Q}_\sigma$ enjoys the same properties as Q_σ, since it is obtained by a slight perturbation from it. The operator $\tilde{L}_\varepsilon$ is a linear second order operator whose coefficients are supported in $[v_\varepsilon, v_\varepsilon + \ln 4] \times S^1$ and are bounded by a constant multiplied for $\varepsilon^{1/2}$, in $\mathcal{C}^\infty$ topology, where partial derivatives are computed with respect to the vector fields ∂_u and ∂_v.

As a fact, if we take into account the effect of the change of the normal vector field, we would obtain, applying the result of Appendix B of [6], a similar formula where the coefficients of the corresponding operator $\tilde{L}_\varepsilon$ are bounded by a constant multiplied for ε since

$$\tilde{N}_\varepsilon \cdot N_\varepsilon = 1 + \mathcal{O}(\varepsilon)$$

for $t \in [t_\varepsilon, t_\varepsilon + \ln 2]$. Instead, if we take into account the effect of the change in the parameterization, we would obtain a similar formula where the coefficients of the corresponding operator $\tilde{L}_\varepsilon$ are bounded by a constant multiplied for $\varepsilon^{1/2}$. The estimate of the coefficients of $\tilde{L}_\varepsilon$ follows from these considerations.

In the following of this section we will give the detailed proof of the existence of a family of a minimal graph about M_1 and asymptotic to it. The proof relative to the case where the surface M_1 is replaced by M_2 can be obtained easily from the previous one. We recall that the surfaces M_1 and M_1 are respectively invariant with respect to two different mirror symmetries. A normal graph of the function $w = w(u, v)$ about M_1 (respectively about M_2) inherits the same property of symmetry if w is an even (respectively odd) function with respect to the variable u. Then once the proof of the result concerning M_1 will be completed, to obtain the result about M_2 it will be sufficient to replace even functions with odd functions and to repeat the same arguments.

Now, assume that we are given a function $\varphi \in \mathcal{C}^{2,\alpha}(I_\sigma)$ which is even with respect to u, L^2-orthogonal to $e_{\sigma,0}, e_{\sigma,1}$ and such that

$$\|\varphi\|_{\mathcal{C}^{2,\alpha}} \leqslant k\varepsilon. \tag{49}$$

We define

$$w_\varphi(\cdot,\cdot) := \bar{\mathcal{H}}_{v_\varepsilon,\varphi}(\cdot,\cdot),$$

where $v_\varepsilon = -\frac{1}{2}\ln\varepsilon + \mathcal{O}(1)$ and $\bar{\mathcal{H}}$ is introduced in proposition 9.5.
In order to solve the equation (48), we choose $\mu \in (-2, -1)$ and look for u of the form $u = w_\varphi + g$ where $g \in \mathcal{C}^{2,\alpha}_\mu(I_\sigma \times [v_\varepsilon, +\infty))$. Using Proposition 6.5, we can rephrase this problem as a fixed point problem

$$g = S(\varphi, g) \tag{50}$$

where the nonlinear mapping S which depends on ε and φ is defined by

$$S(\varphi, g) := G_{\varepsilon, v_\varepsilon}\left(\tilde{L}_\varepsilon(w_\varphi + g) - \mathcal{L}_\sigma\, w_\varphi + \tilde{Q}_\varepsilon\,(w_\varphi + g)\right).$$

where the operator $G_{\varepsilon, v_\varepsilon}$ is defined in Proposition 6.5. To prove the existence of a fixed point for (50) we need the following

Lemma 7.2. *Let $\mu \in (-2,-1)$. There exist some constants $c_k > 0$ and $\varepsilon_k > 0$, such that*

$$\|S(\varphi, 0)\|_{\mathcal{C}^{2,\alpha}_\mu(I_\sigma \times [v_\varepsilon, +\infty))} \leqslant c_k \varepsilon^{3/2+\mu/2} \tag{51}$$

and, for all $\varepsilon \in (0, \varepsilon_k)$

$$\|S(\varphi, g_2) - S(\varphi, g_1)\|_{\mathcal{C}^{2,\alpha}_\mu(I_\sigma \times [v_\varepsilon, +\infty))} \leqslant \frac{1}{2}\, \|g_2 - g_1\|_{\mathcal{C}^{2,\alpha}_\mu(I_\sigma \times [v_\varepsilon, +\infty))},$$

$$\|S(\varphi_2, g) - S(\varphi_1, g)\|_{\mathcal{C}^{2,\alpha}_\mu(I_\sigma \times [v_\varepsilon, +\infty))} \leqslant c\varepsilon^{\frac{1}{2}+\mu/2}\, \|\varphi_2 - \varphi_1\|_{\mathcal{C}^{2,\alpha}} \tag{52}$$

for all $g, g_1, g_2 \in \mathcal{C}^{2,\alpha}_\mu(I_\sigma \times [v_\varepsilon, +\infty))$ such that $\|g_i\|_{\mathcal{C}^{2,\alpha}_\mu(I_\sigma \times [v_\varepsilon, +\infty))} \leqslant 2c_k\varepsilon^{3/2+\mu/2}$ for all boundary data $\varphi, \varphi_1, \varphi_2 \in \mathcal{C}^{2,\alpha}(S^1)$ satisfying (49).

Proof. We know from proposition 6.5 that $\|G_{\varepsilon, v_\varepsilon}(f)\|_{\mathcal{C}^{2,\alpha}_\mu} \leqslant c\|f\|_{\mathcal{C}^{0,\alpha}_\mu}$, then

$$\|S(\varphi, 0)\|_{\mathcal{C}^{2,\alpha}_\mu(I_\sigma \times [v_\varepsilon, +\infty))} \leqslant c\|\tilde{L}_\varepsilon(w_\varphi) - \mathcal{L}_\sigma\, w_\varphi + \tilde{Q}_\sigma\,(w_\varphi)\,\|_{\mathcal{C}^{0,\alpha}_\mu(I_\sigma \times [v_\varepsilon, +\infty))} \leqslant$$

$$\leqslant c\left(\|\tilde{L}_\varepsilon(w_\varphi)\|_{\mathcal{C}^{0,\alpha}_\mu(I_\sigma \times [v_\varepsilon, \infty))} + \|\mathcal{L}_\sigma\, w_\varphi\|_{\mathcal{C}^{0,\alpha}_\mu(I_\sigma \times [v_\varepsilon, \infty))} + \|\tilde{Q}_\sigma\,(w_\varphi)\,\|_{\mathcal{C}^{0,\alpha}_\mu(I_\sigma \times [v_\varepsilon, \infty))}\right).$$

So we need to find the estimates for the three terms above.
We recall that $|\varphi|_{2,\alpha} \leqslant k\varepsilon$. For all $\mu \in (-2,-1)$, thanks to proposition 9.5 we know that

$$|w_\varphi|_{2,\alpha;[v,v+1]} \leqslant e^{-2(v-v_\varepsilon)}|\varphi|_{2,\alpha} \tag{53}$$

We know that

$$\|w_\varphi\|_{\mathcal{C}^{2,\alpha}_\mu} = \sup_{v \in [v_\varepsilon, +\infty)} e^{-\mu v}|w_\varphi|_{2,\alpha;[v,v+1]} \leqslant \sup_{v \in [v_\varepsilon, \infty)} e^{-\mu v} e^{-2(v-v_\varepsilon)}|\varphi|_{2,\alpha} \leqslant$$

$$\leqslant e^{-\mu v_\varepsilon}|\varphi|_{2,\alpha} \leqslant c_k\varepsilon^{1+\mu/2}.$$

From this inequality and from the estimates of the coefficients of $\tilde{L}_\varepsilon$, it follows that

$$\|\tilde{L}_\varepsilon(w_\varphi)\|_{\mathcal{C}^{0,\alpha}_\mu} \leqslant c\varepsilon^{1/2}\|w_\varphi\|_{\mathcal{C}^{0,\alpha}_\mu} \leqslant c_k\varepsilon^{3/2+\mu/2}.$$

As for $\mathcal{L}_\sigma$, being w_φ an harmonic function we obtain following equality

$$\mathcal{L}_\sigma\, w_\varphi = 2k\, w_\varphi,$$

where $k(u,v) = \sin^2\sigma\, \cos^2(x(u)) + \cos^2\sigma\, \sin^2(y(v)) \leqslant c\varepsilon$, because we have set $\alpha + \beta + \sigma \leqslant \varepsilon$ (see lemma 5.3) and $y_\varepsilon \leqslant y(v) \leqslant \pi$, with $y_\varepsilon = \pi - a_\varepsilon$, where $a_\varepsilon = \mathcal{O}(\sqrt{\varepsilon})$. Therefore, we conclude that

$$\|\mathcal{L}_\sigma\, w_\varphi\|_{\mathcal{C}^{0,\alpha}_\mu(I_\sigma \times [v_\varepsilon, \infty))} \leqslant 2c\varepsilon\|w_\varphi\|_{\mathcal{C}^{0,\alpha}_\mu(I_\sigma \times [v_\varepsilon, \infty))} \leqslant c_k\varepsilon^{2+\mu/2}.$$

The last term is estimated by

$$\|\tilde{Q}_\sigma\,(w_\varphi)\,\|_{\mathcal{C}^{0,\alpha}_\mu(I_\sigma \times [v_\varepsilon, \infty))} \leqslant c_k\varepsilon^{2+\mu/2}.$$

Putting together these estimates we get the first result. The details of other estimates are left to the reader. □

Theorem 7.3. *Let be* $B := \{g \in \mathcal{C}^{2,\alpha}_\mu(I_\sigma \times [v_\varepsilon, +\infty))\,;\, ||g||_{\mathcal{C}^{2,\alpha}_\mu} \leqslant 2c_k \varepsilon^{3/2+\mu/2}\}$. *Then the nonlinear mapping* S *defined above has a unique fixed point* g *in* B.

Proof. The previous lemma shows that, if ε is chosen small enough, the nonlinear mapping S is a contraction mapping from the ball B of radius $2c_k\varepsilon^{3/2+\mu/2}$ in $\mathcal{C}^{2,\alpha}_\mu(I_\sigma \times [v_\varepsilon, V_\sigma])$ into itself. This value comes from the estimate of the norm of $S(\varphi, 0)$. Consequently thanks to the Schauder theorem, S has a unique fixed point v in this ball. □

This argument provides a minimal surface which is close to M_1 (respectively M_2) and has one boundary. This surface, denoted by $S_{t,\gamma,\xi_1,d_t}(\varphi)$, is, close to its boundary, a vertical graph over the annulus $B_{2r_\varepsilon} - B_{r_\varepsilon}$ whose parametrization is given, for $\beta = 0$ by

$$(54)\quad \bar{U}_{t,1}(r,\theta) = -(1+\gamma)\ln(2r) + r\kappa_1\cos\theta - \frac{1+\gamma}{r}\xi_1\cos\theta + d_t + \bar{\mathcal{H}}_{v_\varepsilon,\varphi}(\ln 2r - v_\varepsilon, \theta) + V_t(r,\theta).$$

and, for $\alpha = 0$, by

$$(55)\quad \bar{U}_{t,2}(r,\theta) = -(1+\gamma)\ln(2r) - r\kappa_2\sin\theta - \frac{1+\gamma}{r}\xi_2\sin\theta + d_t + \bar{\mathcal{H}}_{v_\varepsilon,\varphi}(\ln 2r - v_\varepsilon, \theta) + V_t(r,\theta).$$

where $v_\varepsilon = -\frac{1}{2}\ln\varepsilon + \mathcal{O}(1)$. The boundary of the surface corresponds to $r_\varepsilon = \frac{1}{2\sqrt{\varepsilon}}$. The function V_t depends non linearly on ε, ϕ. It satisfies $\|V_t(\varepsilon,\phi_i)(r_\varepsilon\cdot,\cdot)\|_{\mathcal{C}^{2,\alpha}(\bar{B}_2 - B_1)} \leqslant c\varepsilon$ and

$$(56)\qquad \|\bar{V}_t(\varepsilon,\phi)(r_\varepsilon\cdot,\cdot) - \bar{V}_t(\varepsilon,\phi')(r_\varepsilon\cdot,\cdot)\|_{\mathcal{C}^{2,\alpha}(\bar{B}_2 - B_1)} \leqslant c\varepsilon^{1/2}\|\phi - \phi'\|_{\mathcal{C}^{2,\alpha}}.$$

This estimate follows from

$$\|\bar{V}_t(\varepsilon,\phi)(r_\varepsilon\cdot,\cdot) - \bar{V}_t(\varepsilon,\phi')(r_\varepsilon\cdot,\cdot)\|_{\mathcal{C}^{2,\alpha}(\bar{B}_2 - B_1)} \leqslant e^{\mu v_\varepsilon}\|S(\phi,\bar{V}_t) - S(\phi',\bar{V}_t)\|_{\mathcal{C}^{2,\alpha}_\mu(I_\sigma\times[v_\varepsilon,+\infty))}$$

and the estimate (52).

8. The matching of Cauchy data

In this section we shall complete the proof of theorems 1.1, 1.2 and 1.3.

8.1. **The proof of theorem 1.2.**

The proof of theorem 1.2 is articulated in two distinct parts: the proof of the existence of the family $\mathcal{K}_1$ and of the existence of the family $\mathcal{K}_2$.

We start proving the existence of the family $\mathcal{K}_2$. The proof is based on an analytical gluing procedure. The surfaces in family $\mathcal{K}_2$ are symmetric with respect to the vertical plane $\{x_2 = 0\}$. So all of the surfaces involved in the following proof must enjoy the same mirror symmetry property. We will show how to glue a compact piece of a Costa-Hoffman-Meeks type surface with bent catenoidal end to two halves of the KMR example $\widetilde{M}_{\sigma,\alpha,0}$ along the upper and lower boundaries and to a horizontal periodic flat annulus with a removed disk along the middle boundary. All of the surfaces just mentioned have the desired property of symmetry as well as the surfaces obtained by them by a slight perturbation. We will recall below the necessary results we proved in previous sections.

Using the result of section 7, we can obtain a minimal surface that is a perturbation of half the KMR example $\widetilde{M}_{\sigma,\alpha,0}$ and is asymptotic to it. This surface, denoted by $S_{t,\lambda_t,\xi_t,d_t}(\varphi_t)$, can be parameterized over the annular neighbourhood $B_{2r_\varepsilon} - B_{r_\varepsilon}$, of its boundary as the vertical graph of (see (54))

$$\bar{U}_t(r,\theta) = -(1+\lambda_t)\ln(2r) + r\kappa_t\cos\theta - \frac{(1+\lambda_t)}{r}\xi_t\cos\theta + d_t + \bar{\mathcal{H}}_{v_\varepsilon,\varphi_t}(\ln 2r - v_\varepsilon, \theta) + \mathcal{O}_{C^{2,\alpha}_b}(\varepsilon).$$

This surface will be glued to the Costa-Hoffman-Meeks example along its upper boundary. The surface that will be glued along the lower boundary, denoted by $S_{b,\lambda_b,\xi_b,d_b}(\varphi_b)$, can be parameterized in the annulus $B_{2r_\varepsilon} - B_{r_\varepsilon}$, as the vertical graph of

$$\bar{U}_b(r,\theta) = (1+\lambda_b)\ln(2r) + r\kappa_b\cos\theta - \frac{(1+\lambda_b)}{r}\xi_b\cos\theta + d_b + \bar{\mathcal{H}}_{v_\varepsilon,\varphi_b}(\ln 2r - v_\varepsilon, \theta) + \mathcal{O}_{C_b^{2,\alpha}}(\varepsilon).$$

We remark that the functions $\varphi_t, \varphi_b \in C^{2,\alpha}(S^1)$ are even and orthogonal in the L^2-sense to the constant function and to $\theta \to \cos\theta$.

Using the result of section 4, we can construct a minimal graph $S_m(\varphi_m)$ close to a horizontal periodic flat annulus. It can be parameterized, in the neighbourhood $B_{2r_\varepsilon} - B_{r_\varepsilon}$ of its boundary, as the vertical graph of

$$\bar{U}_m(r,\theta) = \tilde{H}_{r_\varepsilon,\varphi_m}(r,\theta) + \mathcal{O}_{C_b^{2,\alpha}}(\varepsilon),$$

where $\tilde{H}_{r_\varepsilon,\varphi_m}$ is the harmonic extension of $\varphi_m \in C^{2,\alpha}(S^1)$ which is an even function orthogonal in the L^2-sense to the constant function.

Thanks to the result of section 3, we can construct a minimal surface $\bar{M}^T_{k,\varepsilon}(\varepsilon/2, \Psi)$, with $\Psi = (\psi_t, \psi_b, \psi_m)$, which is close to a truncated genus k Costa-Hoffman-Meeks surface and has three boundaries. The functions $\psi_t, \psi_b, \psi_m \in C^{2,\alpha}(S^1)$ are even. Moreover ψ_t, ψ_b are L^2-orthogonal to the constant function and to $\theta \to \cos\theta$ and ψ_m is orthogonal to the constant function.

The surface $\bar{M}^T_{k,\varepsilon}(\varepsilon/2, \Psi)$ is, close to its upper and lower boundary, a vertical graph over the annulus $B_{r_\varepsilon} - B_{r_\varepsilon/2}$, whose parametrization is, respectively, given by

$$U_t(r,\theta) = \sigma_t - \ln(2r) - \frac{\varepsilon}{2} r\cos\theta + H_{\psi_t}(s_\varepsilon - \ln 2r, \theta) + \mathcal{O}_{C_b^{2,\alpha}}(\varepsilon),$$

$$U_b(r,\theta) = -\sigma_b + \ln(2r) - \frac{\varepsilon}{2} r\cos\theta + H_{\psi_b}(s_\varepsilon - \ln 2r, \theta) + \mathcal{O}_{C_b^{2,\alpha}}(\varepsilon),$$

where $s_\varepsilon = -\frac{1}{2}\ln\varepsilon$. Nearby the middle boundary the surface is a vertical graph whose parametrization is

$$U_m(r,\theta) = \tilde{H}_{\rho_\varepsilon,\psi_m}\left(\frac{1}{r}, \theta\right) + \mathcal{O}_{C_b^{2,\alpha}}(\varepsilon).$$

The functions $\mathcal{O}_{C_b^{2,\alpha}}(\varepsilon)$ replace the functions $V_t, V_b, V_m, \bar{V}_t, \bar{V}_b, \bar{V}_m$ that appear at the end of sections 4, 3 and 7. They depend nonlinearly on the different parameters and boundary data functions but they are bounded by a constant times ε in $C_b^{2,\alpha}$ topology, where partial derivatives are taken with respect to the vector fields $r\partial_r$ and ∂_θ. We assume that the parameters and the boundary functions are chosen so that

$$(57)\quad |\lambda_t|+|\lambda_b|+|-\lambda_t\ln\left(\varepsilon^{-1/2}\right)+\eta_t|+|\lambda_b\ln\left(\varepsilon^{-1/2}\right)+\eta_b|+\varepsilon^{-1/2}/2|\kappa_t+\varepsilon/2|+\varepsilon^{-1/2}/2|\kappa_b+\varepsilon/2|+$$
$$2\varepsilon^{1/2}(|(1+\lambda_t)\xi_t| + |(1+\lambda_b)\xi_b|) + \|\varphi_t\|_{C^{2,\alpha}(S^1)} + \|\varphi_b\|_{C^{2,\alpha}(S^1)} + \|\varphi_m\|_{C^{2,\alpha}(S^1)} +$$
$$+\|\psi_t\|_{C^{2,\alpha}(S^1)} + \|\psi_b\|_{C^{2,\alpha}(S^1)} + \|\psi_m\|_{C^{2,\alpha}(S^1)} \leqslant k\varepsilon,$$

where $\eta_t = d_t - \sigma_t$, $\eta_b = d_b + \sigma_b$. where the constant $k > 0$ is fixed large enough. It remains to show that, for all ε small enough, it is possible to choose the parameters and boundary functions in such a way that the surface

$$S_{t,\lambda_t,\xi_t,d_t}(\varphi_t) \cup S_{b,\lambda_b,\xi_b,d_b}(\varphi_b) \cup S_m(\varphi_m) \cup M^T_k(\varepsilon/2, \Psi)$$

is a C^1 surface across the boundaries of the different summands. Regularity theory will then ensure that this surface is in fact smooth and by construction it has the desired properties. This will therefore complete the proof of the existence of the family of examples denoted by $\mathcal{K}_1$.

It is necessary to fulfill the following system of equations

$$\begin{cases} U_t(r_\varepsilon,\cdot) = \bar{U}_t(r_\varepsilon,\cdot) \\ U_b(r_\varepsilon,\cdot) = \bar{U}_b(r_\varepsilon,\cdot) \\ U_m(r_\varepsilon,\cdot) = \bar{U}_m(r_\varepsilon,\cdot) \\ \partial_r U_b(r_\varepsilon,\cdot) = \partial_r \bar{U}_b(r_\varepsilon,\cdot) \\ \partial_r U_t(r_\varepsilon,\cdot) = \partial_r \bar{U}_t(r_\varepsilon,\cdot) \\ \partial_r U_m(r_\varepsilon,\cdot) = \partial_r \bar{U}_m(r_\varepsilon,\cdot) \end{cases}$$

on S^1. The first three equations lead to the system

$$\begin{cases} -\lambda_t \ln(2r_\varepsilon) + \eta_t - \left((1+\lambda_t)\frac{\xi_t}{r_\varepsilon}\right)\cos\theta + r_\varepsilon(\kappa_t + \frac{\varepsilon}{2})\cos\theta + \varphi_t - \psi_t = \mathcal{O}_{C^{2,\alpha}_b}(\varepsilon) \\ \lambda_b \ln(2r_\varepsilon) + \eta_b - \left((1+\lambda_b)\frac{\xi_b}{r_\varepsilon}\right)\cos\theta + r_\varepsilon(\kappa_b + \frac{\varepsilon}{2})\cos\theta + \varphi_b - \psi_b = \mathcal{O}_{C^{2,\alpha}_b}(\varepsilon) \\ \varphi_m - \psi_m = \mathcal{O}_{C^{2,\alpha}_b}(\varepsilon). \end{cases} \tag{58}$$

The last three equations give the system

$$\begin{cases} -\lambda_t + \left((1+\lambda_t)\frac{\xi_t}{r_\varepsilon}\right)\cos\theta + r_\varepsilon(\kappa_t + \frac{\varepsilon}{2})\cos\theta - \partial_\theta(\varphi_t + \psi_t) = \mathcal{O}_{C^{1,\alpha}_b}(\varepsilon) \\ \lambda_b + \left((1+\lambda_b)\frac{\xi_b}{r_\varepsilon}\right)\cos\theta + r_\varepsilon(\kappa_b + \frac{\varepsilon}{2})\cos\theta - \partial_\theta(\varphi_b + \psi_b) = \mathcal{O}_{C^{1,\alpha}_b}(\varepsilon) \\ \partial_\theta(\varphi_m + \psi_m) = \mathcal{O}_{C^{1,\alpha}_b}(\varepsilon). \end{cases} \tag{59}$$

To obtain this system we applied the results of lemma 9.6 and 9.7. Here, the functions $\mathcal{O}_{C^{l,\alpha}}(\varepsilon)$ in the above expansions depend nonlinearly on the different parameters and boundary data functions but they are bounded by a constant times ε in $C^{l,\alpha}$ topology. The projection of the first two equations of each system over the L^2-orthogonal complement of $\mathrm{Span}\{1, \cos\theta\}$ and the remaining two equations gives the system

$$\begin{cases} \varphi_t - \psi_t = \mathcal{O}_{C^{2,\alpha}_b}(\varepsilon) \\ \varphi_b - \psi_b = \mathcal{O}_{C^{2,\alpha}_b}(\varepsilon) \\ \varphi_m - \psi_m = \mathcal{O}_{C^{2,\alpha}_b}(\varepsilon) \\ \partial_\theta\varphi_t + \partial_\theta\psi_t = \mathcal{O}_{C^{1,\alpha}_b}(\varepsilon) \\ \partial_\theta\varphi_b + \partial_\theta\psi_b = \mathcal{O}_{C^{1,\alpha}_b}(\varepsilon) \\ \partial_\theta\varphi_m + \partial_\theta\psi_m = \mathcal{O}_{C^{1,\alpha}_b}(\varepsilon). \end{cases} \tag{60}$$

Lemma 8.1. *The operator h defined by*

$$\begin{aligned} C^{2,\alpha}(S^1) &\to C^{1,\alpha}(S^1) \\ \varphi &\to \partial_\theta\varphi \end{aligned}$$

acting on functions that are orthogonal to the constant function in the L^2-sense and are even, is invertible.

Proof ([4]). We observe that if we decompose $\varphi = \sum_{j\geqslant 1} \varphi_j \cos(j\theta)$, then

$$h(\varphi) = -\sum_{j\geqslant 1} j\varphi_j \sin(j\theta),$$

is clearly invertible from $H^1(S^1)$ into $L^2(S^1)$. Now elliptic regularity theory implies that this is also the case when this operator is defined between Hölder spaces. $\square$

Using this result, the system (60) can be rewritten as

$$(\varphi_t, \varphi_b, \varphi_m, \psi_t, \psi_b, \psi_m) = \mathcal{O}_{C^{2,\alpha}}(\varepsilon). \tag{61}$$

Recall that the right hand side depends nonlinearly on $\varphi_t, \varphi_b, \varphi_m, \psi_t, \psi_b, \psi_m$ and also on the parameters $\lambda_t, \lambda_b, \eta_t, \eta_b, \xi_t, \xi_b, \kappa_t, \kappa_b$. We look at this equation as a fixed point problem and fix k large enough. Thanks to estimates (56), (19),(20), (16) and (17), we can use a fixed point theorem for contracting mappings in the ball of radius $k\varepsilon$ in $(C^{2,\alpha}(S^1))^6$ to obtain, for all ε small enough, a solution $(\varphi_t, \varphi_b, \varphi_m, \psi_t, \psi_b, \psi_m)$ of (61). This solution being obtained a fixed point for contraction mapping and the right hand side of (61) being continuous with respect to all data, we see that this fixed point $(\varphi_t, \varphi_b, \varphi_m, \psi_t, \psi_b, \psi_m)$ depends continuously (and in fact smoothly) on the parameters $\lambda_t, \lambda_b, \eta_t, \eta_b, \xi_t, \xi_b, \kappa_t, \kappa_b$. Inserting the founded solution into (58) and (59), we see that it remains to solve a system of the form

$$\left\{\begin{array}{c} -\lambda_t \ln(2r_\varepsilon) + \eta_t + \Big(-(1+\lambda_t)\frac{\xi_t}{r_\varepsilon} + r_\varepsilon(\kappa_t + \frac{\varepsilon}{2})\Big)\cos\theta = \mathcal{O}(\varepsilon), \\ \lambda_b \ln(2r_\varepsilon) + \eta_b + \Big(-(1+\lambda_b)\frac{\xi_b}{r_\varepsilon} + r_\varepsilon(\kappa_b + \frac{\varepsilon}{2})\Big)\cos\theta = \mathcal{O}(\varepsilon), \\ -\lambda_t + \Big((1+\lambda_t)\frac{\xi_t}{r_\varepsilon} + r_\varepsilon(\kappa_t + \frac{\varepsilon}{2})\Big)\cos\theta = \mathcal{O}(\varepsilon), \\ \lambda_b + \Big((1+\lambda_b)\frac{\xi_b}{r_\varepsilon} + r_\varepsilon(\kappa_b + \frac{\varepsilon}{2})\Big)\cos\theta = \mathcal{O}(\varepsilon). \end{array}\right. \tag{62}$$

where this time, the right hand sides only depend nonlinearly on $\lambda_t, \lambda_b, \eta_t, \eta_b, \xi_t, \xi_b, \kappa_t, \kappa_b$. There are eight equations that are obtained by projecting this system over the constant function and the function $\theta \to \cos\theta$. If we set

$$(\bar\eta_t, \bar\eta_b) = (-\lambda_t \ln(2r_\varepsilon) + \eta_t, \lambda_b \ln(2r_\varepsilon) + \eta_b),$$

$$(\bar\xi_t, \bar\xi_b) = r_\varepsilon^{-1}((1+\lambda_t)\xi_t, (1+\lambda_b)\xi_b), \quad (\bar\kappa_t, \bar\kappa_b) = r_\varepsilon(\kappa_t, \kappa_b),$$

the previous system can be rewritten as

$$(\lambda_t, \lambda_b, \bar\xi_t, \bar\xi_b, \bar\eta_t, \bar\eta_b, \bar\kappa_t, \bar\kappa_b) = \mathcal{O}(\varepsilon). \tag{63}$$

This time, provided k has been fixed large enough, we can use Leray-Schäuder fixed point theorem in the ball of radius $k\varepsilon$ in $\mathbb{R}^8$ to solve (63), for all ε small enough. This provides a set of parameters and a set of boundary data such that (58) and (59) hold. Equivalently we have proven the existence of a solution of systems (58) and (59). So the proof of the first part of theorem 1.2 is complete.

The proof of the second part of theorem 1.2 uses the same arguments seen above. So we will omit most of the details. Our aim is showing the existence of the family of surfaces denoted by $\mathcal{K}_1$. The surfaces in the family $\mathcal{K}_1$ are symmetric with respect to the vertical plane $\{x_1 = 0\}$. It is important to observe that in the proof, the KMR example which we deal with, is obtained by slight perturbation of $\widetilde{M}_{\sigma,0,\beta}$. The symmetry properties of this surface differ by the ones of the surface close to $\widetilde{M}_{\sigma,\alpha,0}$ involved in the gluing procedure described above. In particular $\widetilde{M}_{\sigma,0,\beta}$ is symmetric with respect to the vertical plane $\{x_1 = 0\}$. We recall that the Costa-Hoffman-Meeks type surface involved in the first gluing procedure enjoys a mirror symmetry with respect to the vertical plane $\{x_2 = 0\}$. Then it is not appropriate for the gluing with a KMR example of the type described above. To obtain the Costa-Hoffman-Meeks type surface we need, it is sufficient to rotate counterclockwise by $\pi/2$ with respect the vertical axis x_3 the Costa-Hoffman-Meeks surface with bent catenoidal ends described in section 3. The surface obtained in this way enjoys the mirror symmetry with respect to the vertical plane $\{x_1 = 0\}$. In the parametrizations of the top and bottom ends the cosinus function is replaced by the sinus function, that is:

$$U_t(r,\theta) = \sigma_t - \ln(2r) - \frac{\varepsilon}{2} r \sin\theta + H_{\psi_t}(s_\varepsilon - \ln(2r), \theta) + \mathcal{O}_{C^{2,\alpha}_b}(\varepsilon),$$
$$U_b(r,\theta) = -\sigma_b + \ln(2r) - \frac{\varepsilon}{2} r \sin\theta + H_{\psi_b}(s_\varepsilon - \ln(2r), \theta) + \mathcal{O}_{C^{2,\alpha}_b}(\varepsilon),$$

where $s_\varepsilon = -\frac{1}{2}\ln\varepsilon$ and $(r,\theta) \in B_{r_\varepsilon} - B_{r_\varepsilon/2}$. As for the planar middle end, the form of its parametrization remains unchanged. We refer to the first part of the proof for its expression. Another important remark to do, concerns the properties of the Dirichlet boundary data ψ_t, ψ_b, ψ_m. If before it was requested that they were even functions and orthogonal (only ψ_t, ψ_b) to the constant function and to $\theta \to \cos\theta$ to preserve the mirror symmetry property with respect to the vertical plane $\{x_1 = 0\}$, now as consequence of the above observations they must be odd functions and ψ_t, ψ_b must be orthogonal to the constant function and to $\theta \to \sin\theta$. It is clear that all the results showed in section 3 continue to hold.

Now we give the expressions of the parametrizations of the surface $S_{t,\lambda_t,\xi_t,d_t}(\varphi_t)$, the minimal surface obtained by perturbation from the KMR example $\widetilde{M}_{\sigma,0,\beta}$ and asymptotic to it. This surface can be parameterized in the neighbourhood $B_{2r_\varepsilon} - B_{r_\varepsilon}$, as the vertical graph of

$$\bar{U}_t = -(1+\lambda_t)\ln(2r) + r\kappa_t \sin\theta - \frac{(1+\lambda_t)}{r}\xi_t \sin\theta + d_t + \bar{\mathcal{H}}_{v_\varepsilon,\varphi_t}(\ln 2r - v_\varepsilon, \theta) + \mathcal{O}_{C^{2,\alpha}_b}(\varepsilon).$$

The parametrization of the surface that we will glue to the Costa-Hoffman-Meeks type surface along its lower boundary is given by

$$\bar{U}_b(r,\theta) = (1+\lambda_b)\ln(2r) - \xi_b \frac{(1+\lambda_b)}{r}\sin\theta + d_b + \bar{\mathcal{H}}_{v_\varepsilon,\varphi_b}(\ln 2r - v_\varepsilon, \theta) + \mathcal{O}_{C^{2,\alpha}_b}(\varepsilon),$$

where $(r,\theta) \in B_{2r_\varepsilon} - B_{r_\varepsilon}$.

To prove the theorem it is necessary to show the existence of a solution of the following system of equations

$$\begin{cases} U_t(r_\varepsilon,\cdot) = \bar{U}_t(r_\varepsilon,\cdot) \\ U_b(r_\varepsilon,\cdot) = \bar{U}_b(r_\varepsilon,\cdot) \\ U_m(r_\varepsilon,\cdot) = \bar{U}_m(r_\varepsilon,\cdot) \\ \partial_r U_b(r_\varepsilon,\cdot) = \partial_r \bar{U}_b(r_\varepsilon,\cdot) \\ \partial_r U_t(r_\varepsilon,\cdot) = \partial_r \bar{U}_t(r_\varepsilon,\cdot) \\ \partial_r U_m(r_\varepsilon,\cdot) = \partial_r \bar{U}_m(r_\varepsilon,\cdot) \end{cases}$$

on S^1, under the assumption (57) for the parameters and the boundary functions. It is clear that the proof of the existence of a solution of this system is based on the same arguments seen before. We remark that the role played before by the functions $\theta \to \cos(i\theta)$, now is played by the functions $\theta \to \sin(i\theta)$. That completes the proof of theorem 1.2.

8.2. **The proof of theorem 1.1.** To proof the theorem 1.1 we will glue a compact piece of the surface $M^T_k(\xi)$, with $\xi = 0$, described in section 3 to two halves of a Scherk type surface along the upper and lower boundary and to a horizontal periodic flat annulus along the middle boundary. The construction of these surfaces is showed in section 4. In particular we showed the existence of a minimal graph close to half a Scherk type example whose ends have asymptotic directions given by $\cos\theta_1 e_1 + \sin\theta_1 e_3$ and $-\cos\theta_2 e_1 + \sin\theta_2 e_3$. These surfaces in the neighbourhood $B_{2r_\varepsilon} - B_{r_\varepsilon}$ of the boundary, admit the following parametrization

$$\bar{U}_t = d_t - \ln(2r) + \tilde{H}_{r_\varepsilon,\varphi_t}(r,\theta) + \mathcal{O}_{C^{2,\alpha}_b}(\varepsilon),$$
$$\bar{U}_b = d_b + \ln(2r) + \tilde{H}_{r_\varepsilon,\varphi_b}(r,\theta) + \mathcal{O}_{C^{2,\alpha}_b}(\varepsilon),$$

where the Dirichlet boundary data $\varphi_i \in C^{2,\alpha}(S^1)$, for $i = t, b$, are requested to be even and orthogonal to the constant function, w_{φ_i} denotes their harmonic extensions. The other surfaces involved in the gluing procedure have been described in the previous subsection.

The proof is similar to the one given for theorem 1, so we will give only the essential details. Actually to prove the theorem it is necessary to show the existence of a solution of the following system of equations

$$\begin{cases} U_t(r_\varepsilon, \cdot) = \bar{U}_t(r_\varepsilon, \cdot) \\ U_b(r_\varepsilon, \cdot) = \bar{U}_b(r_\varepsilon, \cdot) \\ U_m(r_\varepsilon, \cdot) = \bar{U}_m(r_\varepsilon, \cdot) \\ \partial_r U_b(r_\varepsilon, \cdot) = \partial_r \bar{U}_b(r_\varepsilon, \cdot) \\ \partial_r U_t(r_\varepsilon, \cdot) = \partial_r \bar{U}_t(r_\varepsilon, \cdot) \\ \partial_r U_m(r_\varepsilon, \cdot) = \partial_r \bar{U}_m(r_\varepsilon, \cdot) \end{cases}$$

on S^1, under a assumption similar to (57). We refer to subsection 8.1 for the expressions of $U_t, U_b, U_m, \bar{U}_m$. It is necessary to point out that in this subsection we consider the more symmetric example ($\xi = 0$) in the family $(M_k^T(\xi))_\xi$. So it is necessary to replace $\varepsilon/2$ by 0 in the expressions of the functions U_t and U_b of the top and bottom ends.

We want to remark the boundary data for the surfaces we are going to glue together do not satisfy the same hypotheses of orthogonality. All of these functions are orthogonal to the constant function, but only ψ_t, ψ_b are orthogonal to $\theta \to \cos\theta$ too. The functions denoted by $\mathcal{O}_{C_b^{2,\alpha}}(\varepsilon)$ that appear in the expressions of $\bar{U}_i$ and U_i, with $i = t, b, m$, have a Fourier series decomposition containing a term collinear to $\cos\theta$ only if the corresponding boundary data is assumed to be orthogonal only to the constant function. Furthermore the fact that $\xi = 0$ (the catenoidal ends are not bent) implies that functions which parametrize the top and bottom end of $M_k^T(0)$ are orthogonal to $\cos\theta$. In other terms, in difference with the Scherk type surfaces, we are not able to prescribe the coefficients in front of the eigenfunction $\cos\theta$ for the catenoidal ends of $M_k^T(0)$, because in this more symmetric setting are obliged to vanish.

The first three equations lead to the system

$$\begin{cases} \eta_t + \varphi_t - \psi_t = \mathcal{O}_{C_b^{2,\alpha}}(\varepsilon) \\ \eta_b + \varphi_b - \psi_b = \mathcal{O}_{C_b^{2,\alpha}}(\varepsilon) \\ \varphi_m - \psi_m = \mathcal{O}_{C_b^{2,\alpha}}(\varepsilon), \end{cases} \tag{64}$$

where $\eta_t = d_t - \sigma_t$, $\eta_b = d_b + \sigma_b$. The last three equations give the system

$$\begin{cases} \partial_\theta(\varphi_t + \psi_t) = \mathcal{O}_{C_b^{1,\alpha}}(\varepsilon) \\ \partial_\theta(\varphi_b + \psi_b) = \mathcal{O}_{C_b^{1,\alpha}}(\varepsilon) \\ \partial_\theta(\varphi_m + \psi_m) = \mathcal{O}_{C_b^{1,\alpha}}(\varepsilon). \end{cases} \tag{65}$$

Now to complete the proof it is sufficient to use the same arguments of subsection 8.1.

8.3. **The proof of theorem 1.3.** To prove this theorem it is necessary to treat separately the case $k = 0$ and $k \geqslant 1$.

The case k=0. We will glue half a Scherk example with half a KMR example with $\alpha = \beta = 0$. We observe that this surface is symmetric with respect to the $\{x_1 = 0\}$ and $\{x_2 = 0\}$ planes. The Scherk example is symmetric with respect to the $\{x_2 = 0\}$ plane. To preserve this property of symmetry in the surface obtained by the gluing procedure, we will consider the perturbation of $\tilde{M}_{\sigma,0,0}$ which enjoys the same mirror symmetry. That is the surface denoted by $S_{t,\lambda_t,\xi_t,d_t}(\varphi)$ with $\lambda_t = \xi_t = 0$ and $d_t = d$. It can be parametrized in the annulus $B_{2r_\varepsilon} - B_{r_\varepsilon}$ as the vertical graph of

$$\bar{U}(r,\theta) = -\ln(2r) + \bar{d} + \bar{\mathcal{H}}_{v_\varepsilon,\varphi}(\ln 2r - v_\varepsilon, \theta) + \mathcal{O}_{C^{2,\alpha}_b}(\varepsilon).$$

The Scherk example is parametrized as the vertical graph of

$$U(r,\theta) = -\ln(2r) + d + \tilde{H}_{r_\varepsilon,\psi}(r,\theta) + \mathcal{O}_{C^{2,\alpha}_b}(\varepsilon).$$

As for the Dirichlet boundary data, we assume φ to be an even function orthogonal to the constant function and to $\theta \to \cos\theta$, and ψ to be an even function orthogonal to the constant function.

To prove the theorem in the case $k = 0$, it is necessary to show the existence of a solution of the following system of equations

$$\begin{cases} U(r_\varepsilon,\cdot) = \bar{U}(r_\varepsilon,\cdot) \\ \partial_r U(r_\varepsilon,\cdot) = \partial_r \bar{U}(r_\varepsilon,\cdot) \end{cases}$$

on S^1, under appropriate assumptions on the norms of the Dirichlet boundary data and the parameters ξ, d, $\bar{d}$. These equations lead to the system

$$\begin{cases} \eta + \varphi - \psi = \mathcal{O}_{C^{2,\alpha}_b}(\varepsilon) \\ \partial_\theta(\varphi + \psi) = \mathcal{O}_{C^{1,\alpha}_b}(\varepsilon). \end{cases} \tag{66}$$

where $\eta = \bar{d} - d$. Now to complete the proof it is sufficient to use the same arguments of subsection 8.1. The details are omitted.

The case $k \geqslant 1$. The proof of the theorem in this case is similar the proof of theorem 1.1. In fact three of the surfaces we are going to glue are a compact piece of the Costa-Hoffman-Meeks example M_k, half a Scherk type example and a horizontal periodic flat annulus as in the proof of theorem 1.1. The fourth surface is half a KMR example, of the same type of the proof of theorem for $k = 0$. The surfaces are parametrized as vertical graph over $B_{2r_\varepsilon} - B_{r_\varepsilon}$ of the following functions:

$$\bar{U}_b(r,\theta) = \ln(2r) + d_b + \tilde{H}_{r_\varepsilon,\varphi_b}(r,\theta) + \mathcal{O}_{C^{2,\alpha}_b}(\varepsilon),$$

for the Scherk type example,

$$\bar{U}_m(r,\theta) = \tilde{H}_{r_\varepsilon,\varphi_m}(r,\theta) + \mathcal{O}_{C^{2,\alpha}_b}(\varepsilon),$$

for the horizontal periodic flat annulus,

$$\bar{U}_t(r,\theta) = -\ln(2r) + d_t + \bar{\mathcal{H}}_{v_\varepsilon,\varphi_t}(\ln 2r - v_\varepsilon, \theta) + \mathcal{O}_{C^{2,\alpha}_b}(\varepsilon),$$

for the KMR example,

$$U_t(r,\theta) = \sigma_t - \ln(2r) + H_{\psi_t}(s_\varepsilon - \ln(2r), \theta) + \mathcal{O}_{C^{2,\alpha}_b}(\varepsilon),$$

$$U_b(r,\theta) = -\sigma_b + \ln(2r) + H_{\psi_b}(s_\varepsilon - \ln(2r), \theta) + \mathcal{O}_{C^{2,\alpha}_b}(\varepsilon),$$

$$U_m(r,\theta) = \tilde{H}_{\rho_\varepsilon,\varphi_m}(\frac{1}{r}, \theta) + \mathcal{O}_{C^{2,\alpha}_b}(\varepsilon)$$

for the compact piece of the Costa-Hoffmam-Meeks example. The Dirichlet boundary data are requested to be even functions. Functions ψ_t, ψ_b are orthogonal to the constant function and to

$\theta \to \cos\theta$. Functions ψ_m, φ_t, φ_b, φ_m are orthogonal only to the constant function. In this case the system of equations to solve is:

$$\left\{\begin{array}{c} \eta_t + \varphi_t - \psi_t = \mathcal{O}_{C^{2,\alpha}_b}(\varepsilon) \\ \eta_b + \varphi_b - \psi_b = \mathcal{O}_{C^{2,\alpha}_b}(\varepsilon) \\ \varphi_m - \psi_m = \mathcal{O}_{C^{2,\alpha}_b}(\varepsilon) \\ \partial_\theta(\varphi_t + \psi_t) = \mathcal{O}_{C^{1,\alpha}_b}(\varepsilon) \\ \partial_\theta(\varphi_b + \psi_b) = \mathcal{O}_{C^{1,\alpha}_b}(\varepsilon) \\ \partial_\theta(\varphi_m + \psi_m) = \mathcal{O}_{C^{1,\alpha}_b}(\varepsilon), \end{array}\right. \tag{67}$$

where $\eta_t = d_t - \sigma_t$, $\eta_b = d_b + \sigma_b$. The details are left to the reader.

9. Appendix A

Definition 9.1. *Given $\ell \in \mathbb{N}$, $\alpha \in (0,1)$ and $\nu \in \mathbb{R}$, the space $\mathcal{C}^{\ell,\alpha}_\nu(B_{\rho_0}(0))$ is defined to be the space of functions in $\mathcal{C}^{\ell,\alpha}_{loc}(B_{\rho_0}(0))$ for which the following norm is finite*

$$\|\rho^{-\nu}\, w\|_{\mathcal{C}^{\ell,\alpha}(B_{\rho_0}(0))}.$$

Now we can state the following result.

Proposition 9.2. *There exists an operator*

$$\tilde{H} : C^{2,\alpha}(S^1) \longrightarrow C^{2,\alpha}_0(S^1 \times [\bar\rho, +\infty)),$$

such that for each even function $\varphi(\theta) \in C^{2,\alpha}(S^1)$, which is L^2-orthogonal to the constant function, then $w_\varphi = \tilde{H}_{\bar\rho,\varphi}$ solves

$$\left\{\begin{array}{lll} \delta w_\varphi = 0 & on & S^1 \times [\bar\rho, +\infty) \\ w_\varphi = \varphi & on & S^1 \times \{\bar\rho\}. \end{array}\right.$$

Moreover,

$$||\tilde{H}_{\bar\rho,\varphi}||_{C^{2,\alpha}_{-1}(S^1\times[\bar\rho,+\infty))} \leqslant c\,||\varphi||_{C^{2,\alpha}(S^1)}, \tag{68}$$

for some constant $c > 0$.

Remark 9.3. *Following the arguments of the proof below, it is possible to state a similar proposition but with the hypothesis φ odd.*

Proof. We consider the decomposition of the function φ with respect to the basis $\{\cos(i\theta)\}$, that is

$$\varphi = \sum_{i=1}^{\infty} \varphi_i \cos(i\theta).$$

Then the solution w_φ is given by

$$w_\varphi(\rho, \theta) = \sum_{i=1}^{\infty} \left(\frac{\bar\rho}{\rho}\right)^i \varphi_i \cos(i\theta).$$

Since $\frac{\bar\rho}{\rho} \leqslant 1$, then $\left(\frac{\bar\rho}{\rho}\right)^i \leqslant \left(\frac{\bar\rho}{\rho}\right)$, we can conclude that $|w(r,\theta)| \leqslant c\rho^{-1}|\varphi(\theta)|$ and then $||w_\varphi||_{C^{2,\alpha}_{-1}} \leqslant c||\varphi||_{C^{2,\alpha}}$. □

Now we give the statement of an useful result whose proof is contained in [4].

Proposition 9.4. *There exists an operator*

$$H : \mathcal{C}^{2,\alpha}(S^1) \longrightarrow \mathcal{C}^{2,\alpha}_{-2}([0,+\infty) \times S^1),$$

such that for all $\varphi \in \mathcal{C}^{2,\alpha}(S^1)$*, even function and orthogonal to* 1 *and* $\cos\theta$*, in the* L^2*-sense, the function* $w = H_\varphi$ *solves*

$$\left\{ \begin{array}{rcll} (\partial_s^2 + \partial_\theta^2)\, w & = & 0 & \text{in} \quad [0,+\infty) \times S^1 \\ w & = & \varphi & \text{on} \quad \{0\} \times S^1 \end{array} \right.$$

Moreover

$$\|H_\varphi\|_{\mathcal{C}^{2,\alpha}_{-2}} \leqslant c\, \|\varphi\|_{\mathcal{C}^{2,\alpha}},$$

for some constant $c > 0$.

Proposition 9.5. *There exists an operator*

$$\bar{\mathcal{H}}_{v_0} : C^{2,\alpha}(S^1) \longrightarrow C^{2,\alpha}_{\mu}(S^1 \times [v_0,+\infty)),$$

$\mu \in (-2,-1)$*, such that for every function* $\varphi(v) \in C^{2,\alpha}(S^1)$*, which is* L^2*-orthogonal to* $e_{0,i}(u)$ *with* $i = 0,1$ *and even, the function* $w_\varphi = \bar{\mathcal{H}}_{v_0}(\varphi)$ *solves*

$$\left\{ \begin{array}{lll} \partial^2_{uu} w_\varphi + \partial^2_{vv} w_\varphi = 0 & \text{on} & S^1 \times [v_0,+\infty) \\ w_\varphi = \varphi & \text{on} & S^1 \times \{v_0\}. \end{array} \right.$$

Moreover,

$$||\bar{\mathcal{H}}_{v_0}(\varphi)||_{C^{2,\alpha}_{\mu}(S^1 \times [v_0,+\infty])} \leqslant c\, ||\varphi||_{C^{2,\alpha}(S^1)}, \tag{69}$$

for some constant $c > 0$.

Proof. We consider the decomposition of the function φ with respect to the basis $\{e_{0,i}(u)\}$, that is

$$\varphi = \sum_{i=2}^{\infty} \varphi_i e_{0,i}(u).$$

Then the solution w_φ is given by

$$w_\varphi(u,v) = \sum_{i=2}^{\infty} e^{-i(v-v_0)} \varphi_i e_{0,i}(u).$$

We recall that $\mu \in (-2,-1)$ so we have $-i \leqslant \mu$ from which it follows $|w_\varphi|_{2,\alpha;[v,v+1]} \leqslant e^{\mu(v-v_0)}|\varphi|_{2,\alpha}$ and

$$\|w_\varphi\|_{\mathcal{C}^{2,\alpha}_{\mu}} = \sup_{v \in [v_0,\infty]} e^{-\mu v} |w_\varphi|_{2,\alpha;[v,v+1]} \leqslant \sup_{v \in [v_0,\infty]} e^{-\mu v} e^{\mu(v-v_0)} |\varphi|_{2,\alpha} \leqslant e^{-\mu v_0} |\varphi|_{2,\alpha}.$$

□

Lemma 9.6. *Let* $u(r,\theta)$ *be the harmonic extension defined on* $[r_0,+\infty) \times S^1$ *of the even funcion* $\varphi \in \mathcal{C}^{2,\alpha}(S^1)$ *and such that* $u(r_0,\theta) = \varphi(\theta)$*. Then*

$$\partial_\theta u(r,\theta - \pi/2)_{|r=r_0} = -r_0 \partial_r u(r,\theta)_{|r=r_0}.$$

Proof. If $\varphi(\theta) = \sum_{i \geqslant 0} \varphi_i \cos(i\theta)$, then the function u is given by

$$u(r,\theta) = \sum_{i \geqslant 0} \varphi_i \left(\frac{r}{r_0}\right)^i \cos(i\theta).$$

Then

$$\partial_r u(r,\theta) = \sum_{i \geqslant 1} \varphi_i \left(\frac{r}{r_0}\right)^i \frac{i\cos(i\theta)}{r}$$

and

$$\partial_\theta u(r,\theta) = -\sum_{i \geqslant 0} \varphi_i \left(\frac{r}{r_0}\right)^i i\sin(i\theta).$$

Consequently

$$\partial_\theta u(r,\theta - \pi/2) = -\sum_{i \geqslant 0} \varphi_i \left(\frac{r}{r_0}\right)^i i\cos(i\theta)$$

from which lemma follows easily. □

Lemma 9.7. *Let $u(r,\theta)$ be the harmonic extension defined on $[0,r_0] \times S^1$ of the even funcion $\varphi \in \mathcal{C}^{2,\alpha}(S^1)$ and such that $u(r_0,\theta) = \varphi(\theta)$. Then*

$$\partial_\theta u(r,\theta - \pi/2)_{|r=r_0} = r_0 \partial_r u(r,\theta)_{|r=r_0}.$$

Proof. If $\varphi(\theta) = \sum_{i \geqslant 0} \varphi_i \cos(i\theta)$, then the function u is given by

$$u(r,\theta) = \sum_{i \geqslant 0} \varphi_i \left(\frac{r_0}{r}\right)^i \cos(i\theta).$$

Then

$$\partial_r u(r,\theta) = -\sum_{i \geqslant 1} \varphi_i \left(\frac{r_0}{r}\right)^i \frac{i\cos(i\theta)}{r}$$

and

$$\partial_\theta u(r,\theta) = -\sum_{i \geqslant 0} \varphi_i \left(\frac{r_0}{r}\right)^i i\sin(i\theta).$$

Consequently

$$\partial_\theta u(r,\theta - \pi/2) = -\sum_{i \geqslant 0} \varphi_i \left(\frac{r_0}{r}\right)^i i\cos(i\theta)$$

from which lemma follows easily. □

10. APPENDIX B

Proof of proposition 7.1. Let Z be the immersion of the surface $\widetilde{M}_{\sigma,\alpha,\beta}$ and N its normal vector. We want to find the differential equation to which a function f must satisfy such that the surface parametrized by $Z_f = Z + fN$ is minimal. In section 5.2 we parametrized the surface $\widetilde{M}_{\sigma,\alpha,\beta}$ on the cylinder $\mathbb{S}^1 \times \mathbb{R}$. We introduced the map $\mathbf{z}(x,y) : \mathbb{S}^1 \times [0,\pi[\to \bar{\mathbb{C}}$ where x, y denote the sphero-conal coordinates. We start working with the conformal variables p, q defined to be as the real and the imaginary part of $\mathbf{z}$. It holds that

$$|Z_p|^2 = |Z_q|^2 = \Lambda, \quad |N_p|^2 = |N_q|^2 = -K\Lambda,$$
$$\langle N_p, N\rangle = \langle N_q, N\rangle = 0, \quad \langle Z_p, Z_q\rangle = 0, \quad \langle N_p, N_q\rangle = 0,$$
$$\langle N_q, Z_q\rangle = -\langle N_p, Z_p\rangle, \quad \langle N_q, Z_p\rangle = \langle N_p, Z_q\rangle,$$

so

$$\langle N_p, Z_p\rangle = |N_p||Z_p|\cos\gamma_1 = \sqrt{-K}\Lambda\cos\gamma_1,$$
$$\langle N_p, Z_q\rangle = |N_p||Z_q|\cos\gamma_2 = \sqrt{-K}\Lambda\cos\gamma_2.$$

Here K denotes the Gauss curvature, Z_p, Z_q and N_p, N_q denote the partial derivatives of the vectors Z and N, γ_1 is the angle between the vectors N_p and Z_p, γ_2 is the angle between the vectors N_p and Z_q.

The proof of proposition 7.1 is articulated in some lemmas. We denote by E_f, F_f, G_f the coefficiens of the second fundamental form for the surface parametrized by Z_f. The following lemma gives the expression of the area energy functional.

Lemma 10.1.

$$A(f) := \int \sqrt{E_f G_f - F_f^2}\, dp\, dq,$$

with

$$\begin{aligned} E_f G_f - F_f^2 &= \Lambda^2 + \Lambda(f_p^2 + f_q^2) + 2K\Lambda^2 f^2 + 2f(f_q^2 - f_p^2)\sqrt{-K}\Lambda \cos\gamma_1 \\ &\quad -4f f_p f_q \sqrt{-K}\Lambda \cos\gamma_2 - K\Lambda f^2(f_p^2 + f_q^2) + f^4 K^2 \Lambda^2. \end{aligned}$$

Proof. The coefficients of the second fundamental form are:

$$E_f = |\partial_p Z_f|^2 = |Z_p|^2 + f_p^2 + f^2|N_p|^2 + 2f\langle N_p, Z_p\rangle,$$

$$G_f = |\partial_q Z_f|^2 = |Z_q|^2 + f_q^2 + f^2|N_q|^2 + 2f\langle N_q, Z_q\rangle,$$

$$F_f = |\partial_p Z_f \cdot \partial_q Z_f| = f_p f_q + f(\langle Z_p, N_q\rangle + \langle Z_q, N_p\rangle).$$

Then

$$\begin{gathered} E_f G_f = |Z_p|^2|Z_q|^2 + f_p^2|Z_q|^2 + f_q^2|Z_p|^2 + f^2(|N_q|^2|Z_p|^2 + |N_p|^2|Z_q|^2) + \\ f^2(f_p^2|N_q|^2 + f_q^2|N_p|^2) + f^4|N_p|^2|N_q|^2 + 4f^2(\langle N_p Z_p\rangle)(\langle N_q Z_q\rangle) + 2f(f_p^2\langle N_q, Z_q\rangle + f_q^2\langle N_p, Z_p\rangle) + \\ f_p^2 f_q^2 + 2f(\langle N_q, Z_q\rangle|Z_p|^2 + \langle N_p, Z_p\rangle|Z_q|^2) + 2f^3(\langle N_q, Z_q\rangle|Z_p|^2 + \langle N_p, Z_p\rangle|Z_q|^2). \end{gathered}$$

Since $\langle N_q, Z_q\rangle + \langle N_p, Z_p\rangle = 0$ and $|Z_p|^2 = |Z_q|^2$ we can conclude that the last two terms of the previous expression are zero. Since $\langle N_q, Z_p\rangle = \langle N_p, Z_q\rangle$ we have

$$F_f = f_p f_q + 2f\langle N_p, Z_q\rangle.$$

Then

$$F_f^2 = f_p^2 f_q^2 + 4f^2(\langle N_p, Z_q\rangle)^2 + 4f f_p f_q \langle N_p, Z_q\rangle.$$

So the expression of $E_f G_f - F_f^2$ is:

$$\begin{gathered} |Z_p|^2|Z_q|^2 + f_p^2|Z_q|^2 + f_q^2|Z_p|^2 + f^2(|N_q|^2|Z_p|^2 + |N_p|^2|Z_q|^2) + \\ f^2(f_p^2|N_q|^2 + f_q^2|N_p|^2) + f^4|N_p|^2|N_q|^2 + 4f^2(\langle N_p, Z_p)(\langle N_q, Z_q\rangle) + 2f(f_p^2\langle N_q, Z_q\rangle + f_q^2\langle N_p, Z_p\rangle) \\ -4f^2(\langle N_p, Z_q\rangle)^2 - 4f f_p f_q \langle N_p, Z_q\rangle. \end{gathered}$$

Ordering the terms we get:

$$\begin{gathered} |Z_p|^2|Z_q|^2 + f_p^2|Z_q|^2 + f_q^2|Z_p|^2 + f^2(|N_q|^2|Z_p|^2 + |N_p|^2|Z_q|^2) - 4f^2(\langle N_p, Z_q\rangle)^2 \\ +4f^2(\langle N_p, Z_p\rangle)(\langle N_q, Z_q\rangle) + 2f(f_p^2\langle N_q, Z_q\rangle + f_q^2\langle N_p, Z_p\rangle) - 4f f_p f_q \langle N_p, Z_q\rangle + \\ +f^2(f_p^2|N_q|^2 + f_q^2|N_p|^2) + f^4|N_p|^2|N_q|^2. \end{gathered}$$

The expression of $E_f G_f - F_f^2$ becomes:

$$\begin{gathered} \Lambda^2 + \Lambda(f_p^2 + f_q^2) - 2K\Lambda^2 f^2 + 4f^2 K\Lambda^2\left(\cos^2\gamma_1 + \cos^2\gamma_2\right) + \\ +2f(f_q^2 - f_p^2)\sqrt{-K}\Lambda\cos\gamma_1 - 4f f_p f_q\sqrt{-K}\Lambda\cos\gamma_2 - K\Lambda f^2(f_p^2 + f_q^2) + f^4 K^2 \Lambda^2. \end{gathered}$$

Using the relations $\langle N_q, Z_p\rangle = \langle N_p, Z_q\rangle$ and $\langle N_q, Z_q\rangle = -\langle N_p, Z_p\rangle$, it is possible to understand that the relative positions of these vectors are such that $\gamma_2 = \frac{\pi}{2} \pm \gamma_1$. So $\cos^2\gamma_2 = \cos^2(\frac{\pi}{2} \pm \gamma_1) = \sin^2\gamma_1$ and $\cos^2\gamma_1 + \cos^2\gamma_2 = 1$. Then we can write:

$$\Lambda^2 + \Lambda(f_p^2 + f_q^2) + 2K\Lambda^2 f^2 + 2f(f_q^2 - f_p^2)\sqrt{-K}\Lambda\cos\gamma_1$$
$$-4ff_pf_q\sqrt{-K}\Lambda\cos\gamma_2 - K\Lambda f^2(f_p^2 + f_q^2) + f^4K^2\Lambda^2.$$

□

The next lemma completes the proof of the proposition 7.1.

Lemma 10.2. *The surface whose immersion is given by* $Z + fN$, *is minimal if and only if* f *satisfies*

$$\mathcal{L}_\sigma f + Q_\sigma(f) = 0,$$

where $\mathcal{L}_\sigma$ *is the Lamé operator and* Q_σ *is a second order differential operator which satisfies*

$$\|Q_\sigma(f_2) - Q_\sigma(f_1)\|_{\mathcal{C}^{0,\alpha}(I_\sigma\times[v,v+1])} \leqslant c \sup_{i=1,2} \|f_i\|_{\mathcal{C}^{2,\alpha}(I_\sigma\times[v,v+1])} \|f_2 - f_1\|_{\mathcal{C}^{2,\alpha}(I_\sigma\times[v,v+1])}.$$

Proof. The surface parameterized by $Z_f = Z + fN$ is minimal if and only the first variation of $A(f)$ is 0. That is

$$2\,DA_{|f}(g) = \int \frac{1}{\sqrt{(E_fG_f - F_f^2)}_{|f=0}} D_f(E_fG_f - F_f^2)\,(g)\,dp\,dq.$$

Thanks to the previous lemma it holds that

$$\frac{1}{\sqrt{(E_fG_f - F_f^2)}_{|f=0}} D_f(E_fG_f - F_f^2)(g) = \frac{1}{\Lambda}\left(2\Lambda(f_pg_p + f_qg_q) + 4K\Lambda^2 fg + \right.$$
$$+2\sqrt{-K}\Lambda\cos\gamma_1\left[2ff_qg_q + gf_q^2 - 2ff_pg_p - gf_p^2\right] +$$
$$-4\sqrt{-K}\Lambda\cos\gamma_2\left[ff_qg_p + fg_qf_p + gf_pf_q\right] +$$
$$\left.-2K\Lambda\left[fgf_p^2 + f_pg_pf^2 + fgf_q^2 + f_qg_qf^2\right] + 4K^2\Lambda^2f^3g\right).$$

Reordering the summands, we have:

$$\frac{1}{\sqrt{(E_fG_f - F_f^2)}_{|f=0}} D_f(E_fG_f - F_f^2)(g) = 2\left(f_pg_p + f_qg_q + 2K\Lambda fg + \right.$$
$$+\sqrt{-K}\cos\gamma_1\left[2f(f_qg_q - f_pg_p) + g(f_q^2 - f_p^2)\right] +$$
$$-2\sqrt{-K}\cos\gamma_2\left[f(f_qg_p + g_qf_p) + gf_pf_q\right] +$$
$$\left.-K\left[fg(f_p^2 + f_q^2) + f^2(f_pg_p + f_qg_q)\right] + 2K^2\Lambda f^3g\right).$$

In the next computation we can skip the factor 2 in front of the last expression.

$$f_pg_p + f_qg_q + 2K\Lambda fg + Q_1(f, f_p, f_q)g - Q_2(f, f_p, f_q)g_p - Q_3(f, f_p, f_q)g_q = 0,$$

where

$$Q_1(f, f_p, f_q) = -(f_p^2 - f_q^2)\sqrt{-K}\cos\gamma_1 - 2f_pf_q\sqrt{-K}\cos\gamma_2 - Kf(f_p^2 + f_q^2) + 2K^2\Lambda f^3,$$
$$Q_2(f, f_p, f_q) = 2ff_p\sqrt{-K}\cos\gamma_1 + 2ff_q\sqrt{-K}\cos\gamma_2 + Kf^2f_p,$$
$$Q_3(f, f_p, f_q) = -2ff_q\sqrt{-K}\cos\gamma_1 + 2ff_p\sqrt{-K}\cos\gamma_2 + Kf^2f_q.$$

An integration by parts and a change of sign give us the equation:

$$(f_{pp} + f_{qq} - 2K\Lambda f - Q_1(f, f_p, f_q) +$$

$$+P_2(f,f_p,f_q,f_{pp},f_{pq},f_{qq})+P_3(f,f_p,f_q,f_{pp},f_{pq},f_{qq}))\,g=0,$$

where

$$P_2(f,f_p,f_q,f_{pp},f_{pq},f_{qq})=\partial_p Q_2(f,f_p,f_q)$$

and

$$P_3(f,f_p,f_q,f_{pp},f_{pq},f_{qq})=\partial_q Q_3(f,f_p,f_q).$$

That is

$$P_2(f,f_p,f_q,f_{pp},f_{pq},f_{qq})=2(f_p^2+ff_{pp})\sqrt{-K}\cos\gamma_1+2(f_pf_q+ff_{pq})\sqrt{-K}\cos\gamma_2+$$
$$+K(2ff_p^2+f^2f_{pp})+2f(f_p(\sqrt{-K}\cos\gamma_1)_p+f_q(\sqrt{-K}\cos\gamma_2)_p)+f^2f_pK_p$$

and

$$P_3(f,f_p,f_q,f_{pp},f_{pq},f_{qq})=-2(f_q^2+ff_{qq})\sqrt{-K}\cos\gamma_1+2(f_pf_q+ff_{pq})\sqrt{-K}\cos\gamma_2+$$
$$+K(2ff_q^2+f^2f_{qq})+2f(-f_q(\sqrt{-K}\cos\gamma_1)_q+f_p(\sqrt{-K}\cos\gamma_2)_q)+f^2f_qK_q.$$

Now we change the variables, passing from the (p,q) variables to the (u,v) variables. Then we want to understand how above differential equation changes. We recall that p and q are the real and imaginary part of the variable $\mathbf{z}$ of which we know the expression in terms of the spheroconal coordinates x,y (see (28)). It is known that the metric $\bar{g}$ induced on a surface whose immersion Z is given by the Weierstrass representation on a domain of the complex $\mathbf{z}$-plane, can be expressed in terms of the metric $d\bar{s}^2=dp^2+dq^2$, by $\bar{g}=\Lambda(dp^2+dq^2)$, where $\Lambda=|Z_p|^2=|Z_q|^2$. The Laplace-Beltrami operators written with respect to the metrics $d\bar{s}^2$ and $\bar{g}$ are related by:

$$\Delta_{d\bar{s}^2}=\frac{1}{\Lambda}\Delta_{\bar{g}}.$$

That is they differ by the conformal factor $1/\Lambda$. In section 7.1 we observed that the conformal factor related to the change of coordinates $(x,y)\to(u,v)$ is $-K/k$. So the conformal factor due to the change of coordinates $(p,q)\to(u,v)$ is obtained by multiplication of the conformal factors described above. Summarizing it holds that

$$f_{pp}+f_{qq}=\frac{-K\Lambda}{k}(f_{uu}+f_{vv}).$$

So we can write

$$\frac{-K\Lambda}{k}\left(f_{uu}+f_{vv}\right)+2\left(-K\Lambda\right)f+R_1+R_2+R_3=0,$$

where

$$R_1(f,f_u,f_v)=-\frac{-K\Lambda}{k}\left[-(f_u^2-f_v^2)\sqrt{-K}\cos\gamma_1-2f_uf_v\sqrt{-K}\cos\gamma_2-Kf(f_u^2+f_v^2)\right]-2K^2\Lambda f^3$$
$$=\frac{-K\Lambda}{k}\left[(f_u^2-f_v^2)\sqrt{-K}\cos\gamma_1+2f_uf_v\sqrt{-K}\cos\gamma_2+Kf(f_u^2+f_v^2)-2Kkf^3\right]=$$
$$\frac{-K\Lambda}{k}\bar{P}_1(f,f_u,f_v),$$

$$R_2(f,f_u,f_v,f_{uu},f_{uv},f_{vv})=\frac{-K\Lambda}{k}P_2(f,f_u,f_v,f_{uu},f_{uv},f_{vv})$$

and

$$R_3(f,f_u,f_v,f_{uu},f_{uv},f_{vv})=\frac{-K\Lambda}{k}P_3(f,f_u,f_v,f_{uu},f_{uv},f_{vv}).$$

We can write, simplifying the notation:

$$\frac{-K\Lambda}{k}\left[f_{uu}+f_{vv}+2k(u,v)f+\bar{P}_1(f)+P_2(f)+P_3(f)\right]=0.$$

We can recognize the Lamé operator,

$$\mathcal{L}_\sigma f = f_{uu} + f_{vv} + 2(\sin^2\sigma\cos^2 x(u) + \cos^2\sigma\sin^2 y(v))f,$$

then, if we set $Q_\sigma = \bar{P}_1(f) + P_2(f) + P_3(f)$, the equation can be written

$$\mathcal{L}_\sigma f + Q_\sigma(f) = 0.$$

To show the estimate about Q_σ is sufficient to show that all its coefficients are bounded. In particular we will show that the Gauss curvature K and its derivatives K_u, K_v are bounded. We start observing that $\frac{-K}{k(x(u),y(v))}$ is bounded. It is well known that the Gauss curvature has the following expression in terms of the Weierstrass data g, dh:

$$K = -16\left(|g| + \frac{1}{|g|}\right)^{-4}\left|\frac{dg}{g}\right|^2 |dh|^{-2}$$

We recall that $dh = \frac{\mu\, dz}{\sqrt{(z^2+{}^2)(z^2+{}^{-2})}}$. Since $|z^2+\lambda^2||z^2+\lambda^{-2}|$ and $k(x,y) = \sin^2\sigma\,\cos^2 x(u) + \cos^2\sigma\,\sin^2 y(v)$ have the same zeroes, that is the points D, D', D'', D''' given by (22), then $-K/k$ is bounded as well as its derivatives.

We can give an estimate of the derivatives of K and $\sqrt{-K}$. We can write $\sqrt{-K} = \sqrt{k}\sqrt{\frac{-K}{k}}$. From the observations made above it follows that it is sufficient to study the derivatives of $\sqrt{k}$ to show that the derivatives of $\sqrt{-K}$ are bounded.

We recall that

$$l(x) = \sqrt{1 - \sin^2\sigma\sin^2 x} \quad m(y) = \sqrt{1 - \cos^2\sigma\cos^2 y}.$$

From the expression of k, using (29) it is easy to get:

$$\frac{\partial}{\partial u}\sqrt{k} = -\frac{\sin^2\sigma\sin 2x(u)}{2\sqrt{k}}l(x(u)),$$

$$\frac{\partial}{\partial v}\sqrt{k} = \frac{\cos^2\sigma\sin 2y(v)}{2\sqrt{k}}m(y(v)).$$

Then

$$\left|\frac{\partial}{\partial u}\sqrt{k}\right| = \frac{\sin^2\sigma|\sin 2x(u)|l(x(u))}{2\sqrt{\sin^2\sigma\cos^2 x(u) + \cos^2\sigma\sin^2 y(v)}} \leqslant \frac{\sin^2\sigma|\sin 2x(u)|}{2\sin\sigma|\cos x(u)|} \leqslant \sin\sigma,$$

$$\left|\frac{\partial}{\partial v}\sqrt{k}\right| = \frac{\cos^2\sigma|\sin 2y(v)|m(y(v))}{2\sqrt{\sin^2\sigma\cos^2 x(u) + \cos^2\sigma\sin^2 y(v)}} \leqslant \frac{\cos^2\sigma|\sin 2y(v)|}{2\cos\sigma|\sin y(v)|} \leqslant \cos\sigma.$$

That is the derivatives of $\sqrt{k}$ (and consequently the ones of $\sqrt{-K}$) are bounded. That completes the proof.

□

11. APPENDIX C

The differential equation

$$\sin y\,\partial_y\left(\sin y\,\partial_y f\right) - j^2 f + 2\sin^2 y\, f = 0 \tag{70}$$

is a particular case ($l = 1$) of the associated Legendre differential equation, that is given by

$$\sin y\,\partial_y\left(\sin y\,\partial_y f\right) - j^2 f + l(l+1)\sin^2 y\, f = 0,$$

where $l, j \in \mathbb{N}$. The family of the solutions of equation (70) (see [1]) is

$$c_1 P_l^j(\cos y) + c_2 Q_l^j(\cos y),$$

for $l = 1$, where $P_l^j(t)$ and $Q_l^j(t)$ are respectively the associated Legendre functions of first and second kind. If $l = 1$ these functions are defined as follows:

$$P_1^j(t) = \begin{cases} t & \text{if} \quad j = 0 \\ -\sqrt{1-t^2} & \text{if} \quad j = 1 \\ 0, & \text{if} \quad j \geqslant 2, \end{cases}$$

$$Q_1^j(t) = (-1)^j \sqrt{(1-t^2)^j}\, \frac{d^j Q_1^0(t)}{dt^j} \quad \text{and} \quad Q_1^0(t) = \frac{t}{2} \ln\left(\frac{1+t}{1-t}\right) - 1.$$

REFERENCES

[1] M. Abramowitz, I. A. Stegun, *Handbook of mathematical functions with formulas, graphs, and mathematical tables*, National Bureau of Standards Applied Mathematics Series, 55 (1964).

[2] C. J. Costa, *Imersões minimas en* $\mathbb{R}^3$ *de gênero un e curvatura total finita*, PhD thesis, IMPA, Rio de Janeiro, Brasil (1982).

[3] C. J. Costa, *Example of a complete minimal immersion in* $\mathbb{R}^3$ *of genus one and three embedded ends*, Bol. Soc. Brasil. Mat. 15 (1-2), 47-54 (1984).

[4] S. Fakhi, F. Pacard,*Existence result for minimal hypersurfaces with a prescribed finite number of planar ends*, Manuscripta Math., 103 (4), 465-512 (2000).

[5] D. Gilbarg, N. Trudinger, *Elliptic partial differential equations of second order*, Springer-Verlag, Berlin (1998).

[6] L. Hauswirth and F. Pacard, *Higher genus Riemann minimal surfaces*, Invent. Math., 169 (3), 569-620 (2007).

[7] L. Hauswirth and M. Traizet, *The space of embedded doubly-periodic minimal surfaces*, Indiana Univ. Math. J., 51 (5), 1041-1079 (2002).

[8] D. Hoffman and W.H. Meeks III, *A complete embedded minimal surface in* $\mathbb{R}^3$ *with genus one and three ends*, J. Differential Geom., 21 (1), 109-127 (1985).

[9] D. Hoffman and W.H. Meeks III, *The asymptotic behavior of properly embedded minimal surfaces of finite topology*, J. Amer. Math. Soc., 2 (4),667-681 (1989).

[10] D. Hoffman and W.H. Meeks III, *Embedded minimal surfaces of finite topology*, Ann. of Math. (2), 131 (1), 1-34 (1990).

[11] J. K. M. Jansen, *Simple-periodic and non-periodic Lamé functions*, Mathematisch Centrum, Amsterdam (1977).

[12] M. Jleli, *Constant mean curvature hypersurfaces*, PhD Thesis, University Paris 12 (2004).

[13] H. Karcher, *Embedded minimal surfaces derived from Scherk's examples*, Manuscripta Math., 62, 83-114 (1988).

[14] H. Karcher, *Construction of minimal surfaces*, Surveys in Geometry, 1 - 96, University of Tokyo, 1989 and Lecture Notes n. 12, SFB256, Bonn (1989).

[15] T. Kato, *Perturbation theory for linear operators*, Springer-Verlag, Berlin (1980).

[16] N. Korevaar, R. Kusner and B. Solomon, *The Structure of Complete Embedded n. Surfaces with Constant Mean Curvature*, J. Differential Geom., 30 (2), 465-503 (1989).

[17] R. Kusner, R. Mazzeo and D. Pollack, *The moduli space of complete embedded constant mean curvature surfaces*, Geom. Funct. Anal., 6 (1), 120-137 (1996).

[18] F. Martín and V. Ramos Batista, *The embedded syngly periodic Scherk-Costa surfaces*, Math. Ann., 336 (1) 155–189 (2006).

[19] L. Mazet, M. M. Rodríguez and M. Traizet, *Saddle towers with infinitely many ends*, Indiana Univ. Math. J., 56 (6), 2821-2838 (2007).

[20] L. Mazet and M. Traizet, *A quasi-periodic minimal surface*, Preprint.

[21] Meeks III, W. H. and H. Rosenberg, *The Global Theory of Doubly Periodic Minimal Surfaces*, Invent. Math., 97, 351–379 (1989).

[22] R. Melrose, *The Atiyah-Patodi-Singer index theorem*, Research Notes in Mathematics (1993).

[23] F. Morabito, *About a family of deformation of the Costa-Hoffmann-Meeks surfaces*, preprint.

[24] F. Morabito, *Index and nullity of the Gauss map of the Costa-Hoffman-Meeks surfaces*, to appear on Indiana University Mathematics Journal.

[25] S. Nayatani, *Morse index of complete minimal surfaces*, The problem of Plateau, 181-189, World Sci. Publ., River Edge (1992).
[26] S. Nayatani, *Morse index and Gauss maps of complete minimal surfaces in Euclidean 3-space*, Comment. Math. Helv., 68 (4), 511-537 (1993).
[27] J. Pérez, M. M. Rodríguez, M. Traizet, *The classification of doubly periodic minimal tori with parallel ends*, J. Differential Geom., 69 (3), 523-577 (2005).
[28] J. Pérez and A. Ros, *The space of properly embedded minimal surfaces with finite total curvature*, Indiana Univ. Math. J., 45 (1), 177–204 (1996).
[29] V. Ramos Batista, *Singly periodic Costa surfaces*, J. London Math. Soc. 2, 72 (2), 478 - 496 (2005).
[30] M. M. Rodríguez *The space of doubly periodic minimal tori with parallel ends: the standard examples*, Michigan Math. Journal, 55, 103-122 (2007).
[31] W. Rossman, E.C. Thayer and M. Wohlgemuth, *Doubly periodic minimal surfaces*, Experiment. Math., 9 (2), 197–219 (2000).
[32] H.F. Scherk, *Bemerkungen über die kleinste Fläche innerhalb gegebener Grenzen,* J. R. Angew. Math., 13, 185-208 (1835).
[33] F.S. Wei, *Some existence and uniqueness theorems for doubly periodic minimal surfaces*, Invent. Math., 109 (1), 113–136 (1992).

Laurent Hauswirth
Université Paris-Est, Laboratoire d'Analyse et Mathématiques Appliquées, 5 blvd Descartes, 77454 Champs-sur-Marne, FRANCE
E-mail address: `laurent.hauswirth@univ-mlv.fr`

Filippo Morabito
Université Paris-Est, Laboratoire d'Analyse et Mathématiques Appliquées, 5 blvd Descartes, 77454 Champs-sur-Marne, FRANCE
and
Università Roma Tre, Dipartimento di Matematica, Largo S.L. Murialdo 1, 00146 Roma, ITALY
E-mail address: `morabito@mat.uniroma3.it`

M. Magdalena Rodríguez
Universidad Complutense de Madrid, Departamento de Álgebra, P.za de las Ciencias 3, 28040 Madrid, SPAIN
E-mail address: `magdalena@mat.ucm.es`